


Provided for non-commercial research and education use.  
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at ScienceDirect

Cryobiology

journal homepage: [www.elsevier.com/locate/ycryo](http://www.elsevier.com/locate/ycryo)

## The effect of cysteine and glutathione on sperm and oxidative stress parameters of post-thawed bull semen <sup>☆</sup>

Pürhan Barbaros Tuncer <sup>a</sup>, Mustafa Numan Bucak <sup>a,\*</sup>, Serhat Büyükleblebici <sup>a</sup>, Serpil Sarıözkan <sup>b</sup>, Deniz Yeni <sup>c</sup>, Ayşe Eken <sup>d</sup>, Pınar Peker Akalın <sup>e</sup>, Hüseyin Kinet <sup>a</sup>, Fatih Avdatek <sup>c</sup>, A. Fatih Fidan <sup>c</sup>, Mustafa Gündoğan <sup>c</sup>

<sup>a</sup> Tarım ve Köyişleri Bakanlığı, Lalahan Hayvancılık Merkez Araştırma Enstitüsü, Lalahan, Ankara, Turkey

<sup>b</sup> Erciyes Üniversitesi, Safiye Çıkrıkçıoğlu Meslek Yüksekokulu, Kayseri, Turkey

<sup>c</sup> Afyon Kocatepe Üniversitesi, Veteriner Fakültesi, Afyonkarahisar, Turkey

<sup>d</sup> Gülhane Askeri Tıp Akademisi, Etlik, Ankara, Turkey

<sup>e</sup> Tarım ve Köyişleri Bakanlığı, Merkez Veteriner Kontrol ve Araştırma Enstitüsü, Etlik, Ankara, Turkey

### ARTICLE INFO

#### Article history:

Received 17 May 2010

Accepted 20 September 2010

Available online 14 October 2010

#### Keywords:

Bull semen  
Cryopreservation  
Sperm parameters  
Oxidative stress  
Glutathione  
Cysteine

### ABSTRACT

The aim of this study was to determine the effects of antioxidants such as reduced glutathione (GSH) and cysteine in Laiciphose<sup>®</sup> extender on semen parameters, fertilizing ability, lipid peroxidation (LPO) level and glutathione peroxidase (GPx) activity of post-thawed bull semen. Totally 54 ejaculates of three bulls were used in the study. Five groups, namely; GSH (0.5 and 2 mM), cysteine (5 and 10 mM) and control group, were conducted to test the antioxidants in Laiciphose<sup>®</sup>. Insemination doses were processed that each 0.25-mL straw contained  $15 \times 10^6$  sperm. The addition of antioxidants did not present any significant effect on the percentages of post-thaw sperm morphology (acrosome and total abnormalities), subjective, CASA and progressive motilities, as well as sperm motility characteristics (VAP, VSL, VCL, LIN and ALH), compared to the control groups ( $P > 0.05$ ). GSH 0.5 mM ( $55.5 \pm 7.38\%$ ) and cysteine 10 mM ( $48 \pm 5.65\%$ ) led to lower rates of DNA damage, compared to control ( $P < 0.05$ ). As regards to MDA level, cysteine at 10 mM dose gave the highest level ( $4.99 \pm 0.44$  nmol/L) ( $P < 0.001$ ). GPx activity was demonstrated to be higher level upon the addition of 5 mM cysteine when compared to the other groups ( $P < 0.05$ ). With respect to fertility results based on 60-day non-returns, the supplementation of antioxidants did not present significant differences ( $P > 0.05$ ). The results of this study may provide an useful information for the future studies in this area. So, further studies could be suggested to achieve better information in terms of the DNA damage and fertilizing capacity of bull sperm frozen with effective antioxidants.

Crown Copyright © 2010 Published by Elsevier Inc. All rights reserved.

### Introduction

Oxidative stress significantly damages sperm functions due to lipid peroxidation (LPO) induced by reactive oxygen species (ROS). Lipid peroxidation occurs readily in the tissues rich in highly oxidizable polyunsaturated fatty acids (PUFA) [17,10]. Sperm cells contain high concentrations of PUFA, and therefore are highly susceptible to LPO. In addition, every stage of cryopreservation (cooling, freezing and thawing) causes physical and biochemical stress on sperm plasma membrane and structure, which results in a subsequent loss of motility, membrane integrity, fertilizing capability and metabolic changes of sperm [32]. Besides, due to high PUFA

concentration of sperm membrane, mammalian sperm cells may be insufficient in preventing LPO, because of having limited antioxidant system [3,27].

Sperm cells, when frozen and thawed for artificial insemination (AI), have been exposed to oxygen and light radiation which could irreversibly affect sperm functions. Under these conditions, it was stated that supplementation of antioxidants improved the post-thaw motility, viability, membrane integrity and fertility of boar [14], bull [24], ram [30] and goat [2] sperm cells. Cysteine, which has antioxidant capacity, stimulates glutathione synthesis, thus enhancing the maintenance of intracellular glutathione levels and scavenging ROS. Additionally, cysteine has been shown to prevent loss of sperm functions during sperm liquid storage or in the frozen state [22,19]. Glutathione is a large class of antioxidants and has –SH (thio) groups and antioxidant features. Therefore, it is able to react with many ROS directly and is also a co-factor for glutathione peroxidase (GPx) that catalyses the reduction of toxic H<sub>2</sub>O<sub>2</sub> and hydroperoxides [7].

<sup>☆</sup> Statement of funding: This study was supported by Directorate of Lalahan Livestock Central Research Institute, Ankara, Turkey.

\* Corresponding author. Address: Ministry of Agriculture and Rural Affairs, Lalahan Livestock Central Research Institute, Lalahan, Ankara, Turkey.

E-mail address: [mustafanbucak@yahoo.com](mailto:mustafanbucak@yahoo.com) (M.N. Bucak).

In a previous study of bull sperm frozen with Laiciphose extender, the non-return (60–90 days) obtained from field inseminations and in vitro fertilization rates resulted in 60.4% and 83.4–87.1%, respectively [8]. There is a lack of information regarding the effectiveness of commercial Laiciphose extender containing salt buffer and milk powder on sperm functions, LPO level and GPx activity. According to our knowledge, the evaluation of the influence of antioxidants, such as cysteine and GSH, in this extender was not reported. The novelty of this study was to determine the effect of cysteine and GSH, prior to cryopreservation, on semen parameters, fertilizing ability, oxidative stress parameters (MDA and GPx) for frozen-thawed bull semen.

## Materials and methods

### *Animals and semen collection*

In this study, three Holstein bulls (3 and 4 years of age) were housed at the Lalahan Livestock Central Research Institute (Ankara, Turkey), and maintained under uniform feeding and housing conditions. A total number of 60 (20 ejaculates for each bull) ejaculates were collected from these bulls with the aid of an artificial vagina twice a week, according to artificial insemination standard procedures. Ejaculates containing spermatozoa with >80% forward progressive motility and concentrations higher than  $1.0 \times 10^9$  spermatozoa/mL were used in the study. Nineteen ejaculates for each bull met these criteria. The first and second ejaculates from the same bull were combined. The study included nine independent experiments for each bull.

### *Semen processing*

The volume of ejaculates was measured in a conical tube graduated at 0.1 mL intervals and sperm concentration was determined by means of an Accucell photometer (IMV, L'Aigle, France). The sperm motility was estimated by using a phase-contrast microscopy (200 $\times$ ). A extender (Laiciphose: IMV, L'Aigle, France) was used as the base freezing extender. Each ejaculate was split into five equal experimental groups and diluted to a final concentration of  $60 \times 10^6$ /mL spermatozoa with the extender containing cysteine (5 and 10 mM), GSH (0.5 and 2 mM) and no additive (control). Diluted semen samples were loaded into 0.25-mL French straws and cooled down to 4 °C in 2 h, frozen at a programmed rate of  $-3$  °C/min from +4 °C to  $-10$  °C;  $-40$  °C/min from  $-10$  to  $-100$  °C;  $-20$  °C/min from  $-100$  to  $-140$  °C in a digital freezing machine (Digitcool 5300 ZB 250, IMV, France). Thereafter, the straws were plunged into liquid nitrogen. Sperm samples from three bulls were frozen for nine replications. After 24 h, frozen straws were thawed in a 37 °C water bath for 20 s immediately before use.

### *Evaluation of in vitro semen parameters*

Subjective motility was assessed using a phasecontrast microscope (100 $\times$ ), with a warm stage maintained at 37 °C. A wet mount was made using a 5- $\mu$ L drop of semen placed directly on a microscope slide and covered by a coverslip. Sperm motility estimations were performed in three different microscopic fields for each semen sample by the same researcher. The mean of the three successive estimations was recorded as the final motility score. Besides estimating subjective sperm motility, a computer-assisted sperm motility analysis (CASA; IVOS version 12; Hamilton-Thorne Biosciences, MA, USA) was also used to analyze sperm motion characteristics. CASA was set up as follows: phase contrast; frame rate 60 Hz; minimum contrast 80; low and high static size gates 0.10–3.40; low and high intensity gates 0.30–1.70; low and high elongation gates 8–97;

default cell size 5 pixels; default cell intensity 70. Thawed semen was diluted (5 mL semen + 95 mL extender) in a Tris-based extender (without egg yolk and glycerol) and evaluated immediately after dilution. A 4-mL sample of diluted semen was put onto a prewarmed chamber slide (20 mm; Leja 4; Leja Products BV, Holland), and sperm motility characteristics were determined with a 10 $\times$  objective at 37 °C. The following motility values were recorded: motility (%), progressive motility (%), VAP (average path velocity,  $\mu$ m/s), VSL (straight linear velocity,  $\mu$ m/s), VCL (curvilinear velocity,  $\mu$ m/s), ALH (amplitude of lateral head displacement,  $\mu$ m), LIN (linearity index;  $LIN = [VSL/VCL] \times 100$ ). For each evaluation, 10 microscopic fields were analyzed which include at least 300 cells.

For the evaluation of sperm abnormalities, at least three drops of each sample were added to Eppendorf tubes containing 1 mL of Hancock solution [25]. One drop of this mixture was put on a slide and covered with a cover slip. The percentages of acrosome and total abnormalities were determined by counting a total of 400 spermatozoa under phase-contrast microscopy (1000 $\times$  magnification, oil immersion).

The hypo-osmotic swelling test (HOST) was used to evaluate the functional integrity of the sperm membrane. This was performed by incubating 30  $\mu$ L of semen with 300  $\mu$ L of a 100 mOsm hypo-osmotic solution at 37 °C for 60 min. After incubation, 0.2 mL of the mixture was spread with a cover slip on a warm slide. Two hundred sperms with swollen or coiled tails were recorded under phase-contrast microscopy (400 $\times$  magnification) [21].

For assessment of sperm DNA damage, diluted semen samples were centrifuged at 300g for 10 min at 4 °C. Seminal plasma was removed and remaining sperm cells were washed with (Ca<sup>2+</sup> and Mg<sup>2+</sup> free) PBS to yield a concentration of  $1 \times 10^5$  spermatozoa/cm<sup>3</sup> [1]. Sperm DNA damage was investigated using the single cell gel electrophoresis (comet) assay that was generally performed at high alkaline conditions.

### *Embedding of sperm in agarose gel*

Each microscope slide was pre-coated with a layer of 1% normal melting point agarose in PBS and thoroughly dried at room temperature. Next, 100  $\mu$ L of 0.7% low melting point agarose at 37 °C was mixed with 10  $\mu$ L of the cell suspension and dropped on top of the first layer. Slides were allowed to solidify for 5 min at 4 °C in a moist box. The coverslips were removed and the slides were immersed in freshly prepared cold lysis buffer containing 2.5 M NaCl, 100 mM Na<sub>2</sub>-EDTA, 10 mM Tris, 1% Triton X-100 and 40 mM dithiothreitol (pH 10) for 1 h at 4 °C. Then the slides were incubated overnight at 37 °C in 100  $\mu$ g/mL proteinase K and added to the lysis buffer. The slides were removed from the lysis buffer, drained and placed in a horizontal electrophoresis unit filled with fresh alkaline electrophoresis solution, containing 300 mM NaOH and 1 mM EDTA, (pH 13), for 20 min to allow the DNA to unwind. Electrophoresis was performed for 20 min at room temperature at 25 V and was adjusted to 300 mA. Subsequently, the slides were washed with a neutralizing solution of 0.4 M Tris, pH (7.5), in order to remove alkali and detergents. After neutralization the slides were stained with 50  $\mu$ L of 2  $\mu$ L/mL ethidium bromide and covered with a coverslip. All steps were performed under dim light to prevent further DNA damage [16,26]. The images of 100 randomly chosen nuclei were analyzed visually. Observations were made at a magnification of 400 $\times$  using a fluorescent microscope (Olympus, Japan). Each image was classified according to the intensity of the fluorescence in the comet tail, and given a value of 0, 1, 2, 3 or 4 (from undamaged class 0 to maximally damaged class 4), so that the total score of the slide would range from 0 to 400 arbitrary units (AU) [31]. Damage was detected by a tail of fragmented DNA that migrated from the sperm head, causing a 'comet' pattern, whereas whole sperm heads, without a comet, were not considered damaged.

Biochemical assays

MDA level measurement

Supernatant MDA levels were determined in accordance with the method described by Aydin et al. [4]. After the reaction of thiobarbituric acid with MDA, the reaction product was measured spectrometrically. Tetramethoxy propane solution was used as standard. MDA levels of samples were expressed as nmol/mL.

Activity of GPx

GPx activities in samples were measured by the method described by Aydin et al. [4]. The reaction mixture was 50 mmol/L Tris buffer (pH 7.6) containing 1 mmol/L of Na<sub>2</sub>EDTA, 2 mmol/L of reduced glutathione (GSH), 0.2 mmol/L of NADPH, 4 mmol/L of sodium azide and 1000 U of glutathione reductase (GR), 50 µL of samples and 950 µL of reaction mixture were mixed and incubated for 5 min at 37 °C. Then the reaction was initiated with 8 mmol/L of H<sub>2</sub>O<sub>2</sub> and the decrease in NADPH absorbance was followed at 340 nm for 3 min. Enzyme activity was reported in U/mL.

Fertility trials

In fertility trials, the fertilizing ability of spermatozoa was calculated based on the non-return rates. A total of 233 cows were artificially inseminated with frozen samples with antioxidants (a total number of 52, 41, 37, 50 and 53 cows for GSH 0.5 mM, GSH 2 mM, cysteine 5 mM, cysteine 10 mM and control samples, respectively), each comprising the semen of one bull. The effective non-return rates (NNR) were determined at 60 days post-insemination by palpation per rectum.

Statistical analysis

The study included 9 independent experiments for each bull. Results were expressed as the mean ± SEM. Means were analyzed

by analysis of variance (ANOVA), followed by Tukey's post hoc test to determine significant differences in all the parameters between groups using the SPSS/PC computer programme (version 12.0, SPSS, Chicago, IL). Non-return rates were evaluated with chi-square test. Differences with values of *P* < 0.05 were considered to be statistically significant.

Results

Sperm parameters (percentages of motility, acrosome and total abnormalities, HOST and in vivo fertility)

As shown in Table 1, the addition of antioxidants did not present any significant effect on the percentages of post-thaw sperm morphology (acrosome, total abnormalities), subjective, CASA and progressive motilities, as well as sperm motility characteristics (VAP, VCL, VSL, LIN and ALH), compared to the controls. Cysteine at 10 mM dose yielded to slight higher percentage of membrane integrity assessed by HOST than that of the other groups. However, this difference was not statistically significant. GSH 0.5 mM (55.5 ± 7.38%) and cysteine 10 mM (48 ± 5.65%) groups had lower rate of DNA damage compared to control (*P* < 0.05). With respect to fertility results based on 60-day non-returns, the supplementation of antioxidants did not present significant differences (*P* > 0.05).

Analysis of oxidative stress parameters

The influence of antioxidants on LPO level and GPx activity in thawed bull semen is set out in Table 2. Cysteine at 10 mM gave the highest level (4.99 ± 0.44 nmol/mL) of MDA (*P* < 0.001). GPx activity (10.68 ± 0.65) was significantly elevated in the group with cysteine at 5 mM dose when compared to the other groups (*P* < 0.05).

**Table 1**  
Mean (±SE) sperm parameters in frozen–thawed bull semen<sup>a</sup>.

Groups	GSH (±SE)		Cysteine (±SE)		Control (±SE)	P
	0.5 mM	2 mM	5 mM	10 mM		
Subjective motility (%)	48.61 ± 4.93	48.89 ± 4.67	48.61 ± 4.60	48.06 ± 4.69	52.22 ± 2.46	–
CASA motility (%)	46.94 ± 5.24	50.56 ± 4.47	48.94 ± 5.09	51.39 ± 4.82	49.00 ± 3.23	–
Progressive motility (%)	20.11 ± 3.42	22.72 ± 3.07	21.00 ± 3.79	22.06 ± 3.57	21.50 ± 2.28	–
VAP (µm/s)	121.13 ± 3.31	119.09 ± 2.96	120.64 ± 2.96	122.82 ± 2.27	119.69 ± 2.79	–
VSL (µm/s)	101.68 ± 2.91	100.62 ± 2.84	101.36 ± 2.76	104.02 ± 2.52	98.14 ± 2.67	–
VCL (µm/s)	200.16 ± 5.58	196.93 ± 3.86	200.28 ± 4.75	202.02 ± 2.87	202.11 ± 5.76	–
LIN (%)	52.72 ± 0.93	52.89 ± 0.84	52.44 ± 0.82	52.94 ± 0.75	50.83 ± 1.05	–
ALH (µm)	7.68 ± 0.19	7.49 ± 0.19	7.71 ± 0.16	7.67 ± 0.12	7.58 ± 0.24	–
HOST (%)	52.94 ± 4.52	50.33 ± 4.80	47.83 ± 3.90	54.39 ± 4.75	48.22 ± 3.43	–
Acrosome abnormality (%)	1.28 ± 0.30	1.22 ± 0.23	1.33 ± 0.16	0.89 ± 0.17	0.83 ± 0.14	–
Total abnormalities (%)	10.44 ± 0.87	10.28 ± 0.78	10.00 ± 0.56	9.11 ± 0.60	8.83 ± 0.57	–
DNA Damage (AU)	55.5 ± 7.38 <sup>b</sup>	75.5 ± 9.78 <sup>ab</sup>	57 ± 13.3 <sup>ab</sup>	48 ± 5.65 <sup>b</sup>	83.5 ± 6.31 <sup>a</sup>	*
Non-return rates (%)	75.00 (39/52)	60.98 (25/41)	75.68 (28/37)	74.00 (37/50)	69.81 (37/53)	–

<sup>a,b</sup> Different superscripts within the same row demonstrate significant differences (\**P* < 0.05).

–, no significant difference (*P* > 0.05).

<sup>a</sup> The study was replicated nine times.

**Table 2**  
Mean (±SE) malondialdehyde (MDA), and glutathione peroxidase (GPX) activities in frozen–thawed bull semen<sup>a</sup>.

Groups	GSH (±SEM)		Cysteine (±SEM)		Control (±SEM)	P
	0.5 mM	2 mM	5 mM	10 mM		
MDA (nmol/mL)	3.40 ± 0.38 <sup>b</sup>	2.73 ± 0.06 <sup>b</sup>	2.30 ± 0.11 <sup>ab</sup>	4.99 ± 0.44 <sup>c</sup>	1.44 ± 0.08 <sup>a</sup>	***
GPx (U/mL)	8.83 ± 0.99 <sup>ab</sup>	8.28 ± 0.51 <sup>ab</sup>	10.68 ± 0.65 <sup>c</sup>	7.58 ± 0.56 <sup>a</sup>	9.53 ± 0.69 <sup>ab</sup>	*

<sup>a,b,c</sup> Different superscripts within the same row demonstrate significant differences (\**P* < 0.05, \*\*\**P* < 0.001).

<sup>a</sup> The study was replicated nine times.

## Discussion

The increase in sensibility of sperm to oxidative stress and cryodamage affects spermatozoon quality, shortening their life span, resulting in a reduction in sperm motility, viability, antioxidant enzyme activity and fertilizing capacity during extending, freezing and thawing process [18]. Frozen-thawed bull semen are more easily peroxidized than fresh sperm. Additionally, intracellular antioxidant capacity in sperm decreases following freeze-thawing process [28]. In the present study, the addition of GSH and cysteine to the freezing extender did not cause any further improvement in sperm motility parameters. In terms of post-thaw motility, the current results agree with previous studies on boar [15], bull [23,13], and ram semen [9], where no improvement was detected in sperm motility in the presence of GSH. However, motility was not affected by the formation of LPO. This was contradictory to studies in humans [11], mice [5] and horses [6] where deterioration of sperm motility was observed in the presence of free radicals.

Cysteine is a non-essential amino acid and scavenger of free radicals [12]. In this study, the addition of cysteine did not exhibit significant cryoprotective activity on certain sperm functional parameters such as post-thaw motility and morphology, contradictory to the studies performed in bull [24,7], boar [14] and ram [30]. However, cysteine at 5 mM had a beneficial effect on endogenous antioxidant system, improving GPX activity, but not eliminating the MDA production. This finding is in agreement with the result of a study performed on buck [2] sperm in which an increase was observed in the level of GPx activity in the presence of cysteine 15 mM, following the freeze-thawing process.

The integrity of sperm DNA has vital importance to the sperm cell. Some authors suggest that sperm DNA integrity is a more objective marker of sperm function, as opposed to sperm parameters such as motility. DNA damage could be a result of free radical-induced damage [20,29]. In this study, GSH 0.5 mM (55.5 ± 7.38%) and cysteine 10 mM (48 ± 5.65%) gave rise to lower rates of DNA damage, compared to control ( $P < 0.05$ ). It may be stated that, compared to the controls, the use of antioxidants seemed to provide a cryoprotective effect against the DNA damage while reduction of MDA production did not exist during the cryopreservation process.

With respect to fertility results based on 59-day non-returns, the supplementation of antioxidants did not present significant differences of fertility results. This result agrees with the studies which did not observe any improvement in fertility when certain additives were added to the semen extender [24,23].

In conclusion, it may be stated that, compared to the controls, the use of antioxidants in the extender did not eliminate MDA production and improve GPx antioxidant activity during the cryopreservation process, except for GPx in the group with 5 mM cysteine ( $P < 0.05$ ). Sperm motility parameters, morphology and fertility were not influenced by adding of antioxidants. However, DNA integrity was improved when GSH 0.5 mM and cysteine 10 mM were added ( $P < 0.05$ ). The results of this study may provide an useful information for the future studies in this area. So, further studies could be performed to obtain better information in terms of the DNA integrity and fertilizing capacity of bull sperm frozen with effective antioxidants.

## Acknowledgment

This study was supported by the Republic of Turkey, Ministry of Agriculture and Rural Affairs, General Directorate of Agricultural Research (GDAR) (Project No. 07.01.01/02).

## References

- [1] M. Arabi, Bull spermatozoa under mercury stress, *Reprod. Dom. Anim.* 40 (2005) 454–459.
- [2] A. Ateşşahin, M.N. Bucak, P.B. Tuncer, M. Kızıl, Effects of anti-oxidant additives on microscopic and oxidative parameters of Angora goat semen following the freeze-thawing process, *Small Rumin. Res.* 77 (2008) 38–44.
- [3] J.E. Aurich, U. Schönherr, H. Hoppe, C. Aurich, Effect of antioxidants on motility and membrane integrity of chilled-stored stallion semen, *Theriogenology* 48 (1997) 185–192.
- [4] A. Aydın, Z. Arsova-Sarafinovska, A. Sayal, A. Eken, O. Erdem, K. Erten, Y. Özgök, A. Dimovski, Oxidative stress and antioxidant status in non-metastatic prostate cancer and benign prostatic hyperplasia, *Clin. Biochem.* 39 (2006) 176–179.
- [5] G. Baiardi, R.D. Ruiz, M. Cuneo, A.A. Ponce, J.L. Lacuara, L. Vincenti, Differential effects of pharmacologically generated reactive oxygen species upon functional activity of epididymal mouse spermatozoa, *Cancer J. Physiol. Pharmacol.* 75 (1997) 173–178.
- [6] J. Bamber, B.A. Ball, C.G. Gravance, V. Medina, M.C.G. Davies-Morel, The effect of reactive oxygen species on equine sperm motility, viability, acrosomal integrity, mitochondrial potential and membrane lipid peroxidation, *J. Androl.* 21 (2000) 895–902.
- [7] J.F. Bilodeau, S. Blanchette, I.C. Gagnon, M.A. Sirard, Thiols prevent H<sub>2</sub>O<sub>2</sub>-mediated loss of sperm motility in cryopreserved bull semen, *Theriogenology* 56 (2001) 275–286.
- [8] S. Bousseau, J.P. Brillard, B. Marquant-Le Guenne, B. Guérin, A. Camus, M. Lechat, Comparison of bacteriological qualities of various egg yolk sources and the in vitro and in vivo fertilizing potential of bovine semen frozen in egg yolk or lecithin based diluents, *Theriogenology* 50 (1998) 699–706.
- [9] M.N. Bucak, A. Ateşşahin, A. Yüce, Effect of anti-oxidants and oxidative stress parameters on ram semen after the freeze-thawing process, *Small Rumin. Res.* 75 (2008) 128–134.
- [10] S. Chatterjee, E. De Lamirande, C. Gagnon, Cryopreservation alters membrane sulfhydryl status of bull spermatozoa: protection by oxidized glutathione, *Mol. Reprod. Dev.* 60 (2001) 498–506.
- [11] E. De Lamirande, C. Gagnon, Reactive oxygen species and human spermatozoa. I. Effects on the motility of intact spermatozoa and on sperm axonemes, *J. Androl.* 13 (1992) 368–378.
- [12] N. Ercal, P. Treeratphan, P. Lutz, T.C. Hammond, R.H. Mathews, N-acetylcysteine protects Chinese hamster ovary (CHO) cells from lead-induced oxidative stress, *Toxicology* 108 (1996) 57–64.
- [13] R.H. Foote, C.C. Brockett, M.T. Kaproth, Motility and fertility of bull sperm in whole milk extender containing antioxidants, *Anim. Reprod. Sci.* 71 (2002) 13–23.
- [14] H. Funahashi, T. Sano, Selected antioxidants improve the function of extended boar semen stored at 10 °C, *Theriogenology* 63 (2005) 1605–1616.
- [15] J. Gadea, E. Selles, M.A. Marco, P. Coy, C. Matas, R. Romar, S. Ruiz, Decrease in glutathione content in boar sperm after cryopreservation. Effect of the addition of reduced glutathione to the freezing and thawing extenders, *Theriogenology* 62 (2004) 690–701.
- [16] G. Haines, B. Marples, P. Daniel, I. Morris, DNA Damage in human and mouse spermatozoa after in vitro-irradiation assessed by the comet assay, *Adv. Exp. Med. Biol.* 444 (1998) 789–791.
- [17] B. Halliwell, Free radicals and antioxidants: a personal view, *Nutr. Rev.* 52 (1994) 253–265.
- [18] R.H. Hammerstedt, Maintenance of bioenergetic balance in sperm and prevention of lipid peroxidation: a review of the effect on design of storage preservation systems, *Reprod. Fertil.* 5 (1993) 675–690.
- [19] D. Mazor, E. Golan, V. Philip, Red blood cell permeability to thiol compounds following oxidative stress, *Eur. J. Haematol.* 57 (1996) 241–246.
- [20] K.T. Rajesh, K. Doreswamy, B. Shrilatha, M. Muralidhara, Oxidative stress associated DNA damage in testis of mice: induction of abnormal sperms and effects on fertility, *Mutat. Res.* 513 (2002) 103–111.
- [21] S.G. Revel, R.A. Mrode, An osmotic resistance test for bovine semen, *Anim. Reprod. Sci.* 36 (1994) 77–86.
- [22] J. Sagara, K. Miura, S. Bannai, Cysteine uptake and glutathione level in fetal brain cells in primary culture and in suspension, *J. Neurochem.* 61 (1993) 1667–1671.
- [23] S. Sarıözkan, P.B. Tuncer, M.N. Bucak, P.A. Ulutaş, Influence of various antioxidants on microscopic-oxidative stress indicators and fertilizing ability of frozen-thawed bull semen, *Acta Vet. Brno* 78 (2009) 463–469.
- [24] S. Sarıözkan, M.N. Bucak, P.B. Tuncer, P.A. Ulutaş, A. Bilgen, The influence of cysteine and taurine on microscopic-oxidative stress parameters and fertilizing ability of bull semen following cryopreservation, *Cryobiology* 58 (2009) 134–138.
- [25] S. Schafer, A. Holzman, The use of transmigrating and spermac stain to evaluate epididymal cat spermatozoa, *Anim. Reprod. Sci.* 59 (2000) 201–211.
- [26] N.P. Singh, R.E. Stephens, X-ray induced DNA double-strand breaks in human sperm, *Mutagenesis* 13 (1998) 75–79.
- [27] B.T. Storey, Biochemistry of the induction and prevention of lipoperoxidative damage in human spermatozoa, *Mol. Hum. Reprod.* 3 (1997) 203–213.
- [28] G.D. Trincherio, M.A. Affranchino, L.M. Schang, M.T. Beconi, Antioxidant effect of bovine spermatozoa on lipid peroxidation, *Com. Biol.* 8 (1990) 339–350.
- [29] J. Twigg, N. Fulton, E. Gomez, D.S. Irvine, R.J. Aitken, Analysis of the impact of intracellular reactive oxygen species generation on the structural and

- functional integrity of human spermatozoa: lipid peroxidation, DNA fragmentation and effectiveness of antioxidants, *Hum. Reprod.* 13 (1998) 1429–1436.
- [30] O. Uysal, M.N. Bucak, Effects of oxidized glutathione, bovine serum albumin, cysteine and lycopene on the quality of frozen-thawed ram semen, *Acta Vet. Brno* 76 (2007) 383–390.
- [31] F.F. Verit, A. Verit, A. Kocyigit, H. Ciftci, H. Celik, M. Koksak, No increase in sperm DNA damage and seminal oxidative stress in patients with idiopathic infertility, *Arch. Gynecol. Obstet.* 274 (2006) 339–344.
- [32] A.W. Wang, H. Zhang, I. Ikemoto, D.J. Anderson, K.R. Loughlin, Reactive oxygen species generation by seminal cells during cryopreservation, *Urology* 49 (1997) 921–925.