

MULTIVARIABLE CONSTRUCTION OF EXTENDED JACOBI MATRIX POLYNOMIALS

ALİ ÇEVİK

ABSTRACT. The main aim of this paper is to construct a multivariable extension with the help of the extended Jacobi matrix polynomials (EJMPS). Generating matrix functions and recurrence relations satisfied by these multivariable matrix polynomials are derived. Furthermore, general families of multilinear and multilateral generating matrix functions are obtained and their applications are presented.

1. INTRODUCTION

The areas of orthogonal polynomials have many applications in various branches of mathematics and other disciplines. There are many papers in the literature dealing with orthogonal polynomials (see, for example, [3, 14, 24, 29] and the references therein). Recently, matrix orthogonal polynomials have started to become the focus of interest. General theory of matrix valued orthogonal polynomials which have started with the work of M. G. Krein [25, 26] and then have studied by many authors (see [7, 8, 15, 17, 18]) plays an important part in many areas of mathematics just as their scalar counterparts. A good source for matrix polynomials is the book by Gohberg, Lancaster and Rodman [16]. In [1, 5, 2, 4, 9, 6, 10, 12, 20, 21], the classical orthogonal polynomials have been extended to the orthogonal matrix polynomials. Jódar and Cortés introduced and studied the hypergeometric matrix function $F(A, B; C; z)$ and the hypergeometric matrix differential equation in [22] and the explicit closed form general solution of it has been given in [23]. In [6, 11, 10, 12, 19, 20, 27], Jacobi, Chebyshev, Gegenbauer, Laguerre and Hermite matrix polynomials were introduced and various results were given for these matrix polynomials. Furthermore, the authors introduced new extension of Jacobi matrix polynomials in [30]. In [1], authors studied the polynomial $F_n^{(A,B)}(x; a, b, c)$ called extended Jacobi matrix polynomial (EJMP) for parameter matrices A and B whose eigenvalues, z , all satisfy $Re(z) > -1$. For any natural number $n \geq 0$, the n -th

2000 *Mathematics Subject Classification.* 33C45, 33C25, 15A60.

Key words and phrases. Extended Jacobi matrix polynomials; Generating matrix function; Recurrence relation; Multilinear and multilateral generating matrix functions.

©2013 Ilirias Publications, Prishtinë, Kosovë.

Submitted May 13, 2013. Published August 18, 2013.

degree extended Jacobi matrix polynomial $F_n^{(A,B)}(x; a, b, c)$ is defined by

$$F_n^{(A,B)}(x; a, b, c) = \frac{(c(a-b))^n}{n!} F\left(-nI, A+B+(n+1)I; A+I; \frac{x-a}{b-a}\right) (A+I)_n. \quad (1)$$

Also, it is shown that these matrix polynomials have the Rodrigues formula [1]:

$$F_n^{(A,B)}(x; a, b, c) = \frac{(-c)^n}{n!} (x-a)^{-A} (b-x)^{-B} \cdot \frac{d^n}{dx^n} \left\{ (x-a)^{A+nI} (b-x)^{B+nI} \right\}, \quad (c > 0) \quad (2)$$

where A and $B \in \mathbb{C}^{r \times r}$ satisfy

$$\operatorname{Re}(z) > -1 \text{ for } z \in \sigma(A), \operatorname{Re}(\eta) > -1 \text{ for } \eta \in \sigma(B), AB = BA.$$

By comparing the Rodrigues representation for Jacobi matrix polynomials and (2), we have

$$F_n^{(A,B)}(x; a, b, c) = \{c(a-b)\}^n P_n^{(A,B)}\left(\frac{2(x-a)}{a-b} + 1\right) \quad (3)$$

or, equivalently,

$$P_n^{(A,B)}(x) = \{c(a-b)\}^{-n} F_n^{(A,B)}\left(\frac{1}{2}\{a+b+(a-b)x\}; a, b, c\right). \quad (4)$$

The EJMPs $F_n^{(A,B)}(x; a, b, c)$ are orthogonal over the interval (a, b) with respect to the weight function $\omega(x; A, B) = (x-a)^A (b-x)^B$ [1]. In fact, it is hold that

$$\begin{aligned} & \int_a^b (x-a)^A (b-x)^B F_n^{(A,B)}(x; a, b, c) F_m^{(A,B)}(x; a, b, c) dx \\ &= \begin{cases} \frac{c^{2n}}{n!} (b-a)^{A+B+(2n+1)I} \Gamma(A+B+(2n+1)I) \\ \cdot \Gamma^{-1}(A+B+(n+1)I) \Gamma(B+(n+1)I) \\ \cdot \Gamma(A+(n+1)I) \Gamma^{-1}(A+B+2(n+1)I) & , \quad m = n \\ \mathbf{0} & , \quad m \neq n \end{cases} \\ & (m, n \in \mathbb{N}_0 := \mathbb{N} \cup \{0\} = \{0, 1, 2, \dots\}) \end{aligned} \quad (5)$$

where A and $B \in \mathbb{C}^{r \times r}$ satisfy

$$\operatorname{Re}(z) > -1 \text{ for } z \in \sigma(A), \operatorname{Re}(\eta) > -1 \text{ for } \eta \in \sigma(B), AB = BA.$$

In this paper, we construct a multivariable extension of extended Jacobi matrix polynomials and show that these matrix polynomials are orthogonal with respect to weight matrix function. Generating matrix functions are obtained for the multivariable extended Jacobi matrix polynomials (MEJMPs) and with the help of generating matrix function, several recurrence formulas are given for these polynomials. Furthermore, multilinear and multilateral generating matrix functions are derived for MEJMPs and some applications of the results obtained are presented.

Throughout this paper, for a matrix $A \in \mathbb{C}^{r \times r}$, its spectrum is denoted by $\sigma(A)$. The two-norm of A , which will be denoted by $\|A\|$, is defined by

$$\|A\| = \sup_{x \neq 0} \frac{\|Ax\|_2}{\|x\|_2},$$

where, for a vector $y \in \mathbb{C}^r$, $\|y\|_2 = (y^T y)^{1/2}$ is the Euclidean norm of y . I and $\mathbf{0}$ will denote the identity matrix and the null matrix in $\mathbb{C}^{r \times r}$, respectively. We say that a matrix A in $\mathbb{C}^{r \times r}$ is a positive stable if $Re(\lambda) > 0$ for all $\lambda \in \sigma(A)$ where $\sigma(A)$ is the set of all eigenvalues of A . If A_0, A_1, \dots, A_n are elements of $\mathbb{C}^{r \times r}$ and $A_n \neq \mathbf{0}$, then we call

$$P(x) = A_n x^n + A_{n-1} x^{n-1} + \dots + A_1 x + A_0$$

a matrix polynomial of degree n in x . From [22], one can see

$$(P)_n = P(P+I)(P+2I)\dots(P+(n-1)I); \quad n \geq 1; \quad (P)_0 = I. \quad (6)$$

From the relation (6), we see that

$$\frac{(-1)^k}{(n-k)!} I = \frac{(-nI)_k}{n!}; \quad 0 \leq k \leq n. \quad (7)$$

The hypergeometric matrix function $F(A, B; C; z)$ has been given in the form [22]

$$F(A, B; C; z) = \sum_{n=0}^{\infty} \frac{(A)_n (B)_n}{n!} [(C)_n]^{-1} z^n$$

for matrices A, B and C in $\mathbb{C}^{r \times r}$ such that $C + nI$ is invertible for all integer $n \geq 0$ and for $|z| < 1$. For any matrix A in $\mathbb{C}^{r \times r}$, the authors exploited the following relation due to [22]

$$(1-x)^{-A} = \sum_{n=0}^{\infty} \frac{(A)_n}{n!} x^n, \quad |x| < 1. \quad (8)$$

In [13], if $f(z)$ and $g(z)$ are holomorphic functions in an open set Ω of the complex plane, and if A is a matrix in $\mathbb{C}^{r \times r}$ for which $\sigma(A) \subset \Omega$, then

$$f(A)g(A) = g(A)f(A).$$

Hence, if $B \in \mathbb{C}^{r \times r}$ is a matrix for which $\sigma(B) \subset \Omega$ and $AB = BA$, then

$$f(A)g(B) = g(B)f(A).$$

Furthermore, in [12], the reciprocal scalar Gamma function, $\Gamma^{-1}(z) = 1/\Gamma(z)$, is an entire function of the complex variable z . Thus, for any $C \in \mathbb{C}^{r \times r}$, the Riesz-Dunford functional calculus [13] shows that $\Gamma^{-1}(C)$ is well defined and is, indeed, the inverse of $\Gamma(C)$. Hence: if $C \in \mathbb{C}^{r \times r}$ is such that $C + nI$ is invertible for every integer $n \geq 0$, then

$$(C)_n = \Gamma(C + nI)\Gamma^{-1}(C).$$

2. MULTIVARIABLE EXTENSION OF THE EXTENDED JACOBI MATRIX
POLYNOMIALS

A systematic investigation of a multivariable extension of the extended Jacobi matrix polynomials $F_n^{(A,B)}(x; a, b, c)$ is defined by

$$F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) = F_{n_1, \dots, n_s}^{(A_1, \dots, A_s; B_1, \dots, B_s)}(\mathbf{x}) = F_{n_1}^{(A_1, B_1)}(x_1; a_1, b_1, c_1) \dots F_{n_s}^{(A_s, B_s)}(x_s; a_s, b_s, c_s) \quad (9)$$

where $\mathbf{x} = (x_1, \dots, x_s)$, $\mathbf{A} = (A_1, \dots, A_s)$, $\mathbf{B} = (B_1, \dots, B_s)$, A_i and B_i are matrices in $\mathbb{C}^{r \times r}$ satisfying the spectral conditions $Re(z_i) > -1$ for each eigenvalue $z_i \in \sigma(A_i)$ and $Re(\eta_i) > -1$ for each eigenvalue $\eta_i \in \sigma(B_i)$ for $1 \leq i \leq s$ and $\mathbf{n} = n_1 + \dots + n_s$; $n_1, \dots, n_s \in \mathbb{N}_0 := \mathbb{N} \cup \{0\} = \{0, 1, 2, \dots\}$. The multivariable extended Jacobi matrix polynomials $F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x})$ (MEJMPS) are orthogonal with respect to the weight matrix function

$$\begin{aligned} \omega(\mathbf{x}, \mathbf{A}, \mathbf{B}) &= \omega(x_1, \dots, x_s; A_1, \dots, A_s; B_1, \dots, B_s) \\ &= \omega_1(x_1, A_1, B_1) \dots \omega_s(x_s, A_s, B_s) \\ &= (x_1 - a_1)^{A_1} (b_1 - x_1)^{B_1} \dots (x_s - a_s)^{A_s} (b_s - x_s)^{B_s} \end{aligned}$$

over the domain

$$\Omega = \{(x_1, \dots, x_s) : a_i < x_i < b_i ; i = 1, 2, \dots, s\}.$$

In fact, we have by (5)

$$\begin{aligned} &\int_{\Omega} \omega(\mathbf{x}, \mathbf{A}, \mathbf{B}) F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) F_{\mathbf{m}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) d\mathbf{x} \\ &= \int_{a_1}^{b_1} (x_1 - a_1)^{A_1} (b_1 - x_1)^{B_1} F_{n_1}^{(A_1, B_1)}(x_1; a_1, b_1, c_1) F_{m_1}^{(A_1, B_1)}(x_1; a_1, b_1, c_1) dx_1 \times \dots \\ &\quad \times \int_{a_s}^{b_s} (x_s - a_s)^{A_s} (b_s - x_s)^{B_s} F_{n_s}^{(A_s, B_s)}(x_s; a_s, b_s, c_s) F_{m_s}^{(A_s, B_s)}(x_s; a_s, b_s, c_s) dx_s \\ &= \prod_{i=1}^s \left\{ \frac{c_i^{2n_i}}{n_i!} (b_i - a_i)^{A_i + B_i + (2n_i + 1)I} \Gamma(A_i + B_i + (2n_i + 1)I) \Gamma^{-1}(A_i + B_i + (n_i + 1)I) \right. \\ &\quad \cdot \Gamma(B_i + (n_i + 1)I) \Gamma(A_i + (n_i + 1)I) \Gamma^{-1}(A_i + B_i + 2(n_i + 1)I) \left. \right\} \delta_{m_i, n_i} \\ &\quad (m_i, n_i \in \mathbb{N}_0 := \mathbb{N} \cup \{0\} ; i = 1, 2, \dots, s) \end{aligned}$$

where $d\mathbf{x} = dx_1 \dots dx_s$ and these matrices are commutative.

Thus, the following result has been established:

Theorem 1. *The MEJMPS $F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x})$ are orthogonal with respect to the weight matrix function*

$$\omega(\mathbf{x}, \mathbf{A}, \mathbf{B}) = (x_1 - a_1)^{A_1} (b_1 - x_1)^{B_1} \dots (x_s - a_s)^{A_s} (b_s - x_s)^{B_s}$$

over the domain

$$\Omega = \{(x_1, \dots, x_s) : a_i < x_i < b_i ; i = 1, 2, \dots, s\}$$

where A_i and B_i are matrices in $\mathbb{C}^{r \times r}$ satisfying the spectral conditions $Re(z_i) > -1$ for each eigenvalue $z_i \in \sigma(A_i)$ and $Re(\eta_i) > -1$ for each eigenvalue $\eta_i \in \sigma(B_i)$ for $1 \leq i \leq s$ and all matrices are commutative.

3. GENERATING MATRIX FUNCTIONS AND RECURRENCE RELATIONS FOR MEJMPs

In [1], it was shown that the EJMPs are generated by

$$\begin{aligned} & \sum_{n=0}^{\infty} (c(a-b))^{-n} (A+B+I)_n F_n^{(A,B)}(x; a, b, c) [(A+I)_n]^{-1} t^n \\ &= (1-t)^{-(A+B+I)} F\left(\frac{A+B+I}{2}, \frac{A+B+2I}{2}; A+I; \frac{4t(x-a)}{(a-b)(1-t)^2}\right) \end{aligned} \quad (10)$$

where all eigenvalues z of the matrices A and B of the extended Jacobi matrix polynomials $F_n^{(A,B)}(x; a, b, c)$ satisfy the condition $Re(z) > -1$ and $|t| < 1$. On the other hand, other generating function for EJMPs is as follows [1]:

$$\begin{aligned} & \sum_{n=0}^{\infty} (c(a-b))^{-n} F_n^{(A,B)}(x; a, b, c) t^n \\ &= F_4\left(I+B, I+A; I+A, I+B; \frac{(x-a)t}{a-b}, \frac{(x-b)t}{a-b}\right) \end{aligned}$$

where $AB = BA$ and $F_4(A, B; C, D; x, y)$ is the matrix version of the Appell's function of two variables which is defined by

$$\begin{aligned} F_4(A, B; C, D; x, y) &= \sum_{n,k=0}^{\infty} (A)_{n+k} (B)_{n+k} (D)_n^{-1} (C)_k^{-1} \frac{x^k y^n}{k!n!} \\ &(\sqrt{x} + \sqrt{y} < 1) \end{aligned}$$

where $C+nI$ and $D+nI$ are invertible for every integer $n \geq 0$ (see [5]). The other one is

$$\begin{aligned} & \sum_{n=0}^{\infty} (C)_n (D)_n (I+B)_n^{-1} (c(a-b))^{-n} F_n^{(A,B)}(x; a, b, c) (I+A)_n^{-1} t^n \\ &= F_4\left(C, D; I+A, I+B; \frac{(x-a)t}{a-b}, \frac{(x-b)t}{a-b}\right) \end{aligned}$$

where $A+nI$ and $B+nI$ are invertible for every integer $n \geq 0$ [1].

In this section, we obtain generating matrix functions and recurrence relations for MEJMPs $F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x})$.

Using the above expressions, we can give the following results.

Theorem 2. For the MEJMPs $F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x})$, we have

$$\begin{aligned} & \sum_{n_1, \dots, n_s=0}^{\infty} (c(\mathbf{a}-\mathbf{b}))^{-\mathbf{n}} (\mathbf{A}+\mathbf{B}+I)_{\mathbf{n}} [(\mathbf{A}+I)_{\mathbf{n}}]^{-1} F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) t_1^{n_1} \dots t_s^{n_s} \\ &= \prod_{i=1}^s (1-t_i)^{-(A_i+B_i+I)} F\left(\frac{A_i+B_i+I}{2}, \frac{A_i+B_i+2I}{2}; A_i+I; \frac{4t_i(x_i-a_i)}{(a_i-b_i)(1-t_i)^2}\right) \\ &(|t_i| < 1) \end{aligned}$$

or equivalently,

$$\begin{aligned} & \sum_{n_1, \dots, n_s=0}^{\infty} (\mathbf{c}(\mathbf{a} - \mathbf{b}))^{-\mathbf{n}} (\mathbf{A} + \mathbf{B} + I)_{\mathbf{n}} [(\mathbf{A} + I)_{\mathbf{n}}]^{-1} F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})} ((\mathbf{b} - \mathbf{a}) \mathbf{v} + \mathbf{a}) t_1^{n_1} \dots t_s^{n_s} \\ &= \prod_{i=1}^s (1 - t_i)^{-(A_i + B_i + I)} F \left(\frac{A_i + B_i + I}{2}, \frac{A_i + B_i + 2I}{2}; A_i + I; \frac{-4t_i v_i}{(1 - t_i)^2} \right) \\ & \quad (|t_i| < 1) \end{aligned}$$

where A_i and B_i are matrices in $\mathbb{C}^{r \times r}$ satisfying the spectral conditions $\operatorname{Re}(z_i) > -1$ for each eigenvalue $z_i \in \sigma(A_i)$ and $\operatorname{Re}(\eta_i) > -1$ for each eigenvalue $\eta_i \in \sigma(B_i)$ for $1 \leq i \leq s$, all matrices are commutative and

$$\begin{aligned} & \prod_{i=1}^s \left\{ (c_i (a_i - b_i))^{-n_i} (A_i + B_i + I)_{n_i} [(A_i + I)_{n_i}]^{-1} \right\} \\ &= (\mathbf{c}(\mathbf{a} - \mathbf{b}))^{-\mathbf{n}} (\mathbf{A} + \mathbf{B} + I)_{\mathbf{n}} [(\mathbf{A} + I)_{\mathbf{n}}]^{-1}, \\ (\mathbf{b} - \mathbf{a}) \mathbf{v} + \mathbf{a} &= ((b_1 - a_1) v_1 + a_1, \dots, (b_s - a_s) v_s + a_s) \end{aligned}$$

where $A_i + n_i I$ is invertible for every integer $n_i \geq 0$ for $i = 1, 2, \dots, s$.

Theorem 3. The matrix polynomials $F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x})$ are generated by

$$\begin{aligned} & \sum_{n_1, \dots, n_s=0}^{\infty} (\mathbf{c}(\mathbf{a} - \mathbf{b}))^{-\mathbf{n}} F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) t_1^{n_1} \dots t_s^{n_s} \\ &= \prod_{i=1}^s F_4 \left(I + B_i, I + A_i; I + A_i, I + B_i; \frac{(x_i - a_i) t_i}{a_i - b_i}, \frac{(x_i - b_i) t_i}{a_i - b_i} \right) \\ & \quad \left(\sqrt{\frac{(x_i - a_i) t_i}{a_i - b_i}} + \sqrt{\frac{(x_i - b_i) t_i}{a_i - b_i}} < 1; i = 1, 2, \dots, s \right) \end{aligned}$$

where

$$(\mathbf{c}(\mathbf{a} - \mathbf{b}))^{-\mathbf{n}} = \prod_{i=1}^s (c_i (a_i - b_i))^{-n_i}$$

and A_i and B_i are matrices in $\mathbb{C}^{r \times r}$ satisfying the spectral conditions $\operatorname{Re}(z_i) > -1$ for each eigenvalue $z_i \in \sigma(A_i)$ and $\operatorname{Re}(\eta_i) > -1$ for each eigenvalue $\eta_i \in \sigma(B_i)$ for $1 \leq i \leq s$, all matrices are commutative.

Theorem 4. Let $A_i, B_i, C_i, D_i \in \mathbb{C}^{r \times r}$ for $i = 1, 2, \dots, s$. For MEJMPS $F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x})$, we have the following generating matrix function

$$\begin{aligned} & \sum_{n_1, \dots, n_s=0}^{\infty} (\mathbf{C})_{\mathbf{n}} (\mathbf{D})_{\mathbf{n}} (I + \mathbf{B})_{\mathbf{n}}^{-1} (\mathbf{c}(\mathbf{a} - \mathbf{b}))^{-\mathbf{n}} F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) (I + \mathbf{A})_{\mathbf{n}}^{-1} t_1^{n_1} \dots t_s^{n_s} \\ &= \prod_{i=1}^s F_4 \left(C_i, D_i; I + A_i, I + B_i; \frac{(x_i - a_i) t_i}{a_i - b_i}, \frac{(x_i - b_i) t_i}{a_i - b_i} \right) \end{aligned}$$

where $A_i + n_i I$ and $B_i + n_i I$ are invertible for every integer $n_i \geq 0$ for $i = 1, 2, \dots, s$, all matrices are commutative and

$$\begin{aligned} & (\mathbf{C})_{\mathbf{n}} (\mathbf{D})_{\mathbf{n}} (I + \mathbf{B})_{\mathbf{n}}^{-1} (\mathbf{c}(\mathbf{a} - \mathbf{b}))^{-\mathbf{n}} (I + \mathbf{A})_{\mathbf{n}}^{-1} \\ &= \prod_{i=1}^s (C)_{n_i} (D)_{n_i} (I + B_i)_{n_i}^{-1} (c_i (a_i - b_i))^{-n_i} (I + A_i)_{n_i}^{-1}. \end{aligned}$$

Theorem 5. For the MEJMPs $F_{n_1, n_2, \dots, n_s}^{(A_1, \dots, A_s; B_1, \dots, B_s)}(\mathbf{x})$, we have

$$\begin{aligned} & \sum_{n=0}^{\infty} H_n^{(A_1, \dots, A_s; B_1, \dots, B_s)}(x_1, \dots, x_s) t^n \\ &= \prod_{i=1}^s (1-t)^{-(A_i+B_i+I)} F\left(\frac{A_i+B_i+I}{2}, \frac{A_i+B_i+2I}{2}; A_i+I; \frac{4t(x_i-a_i)}{(a_i-b_i)(1-t)^2}\right) \\ & \quad (|t| < 1) \end{aligned}$$

where

$$\begin{aligned} & H_n^{(A_1, \dots, A_s; B_1, \dots, B_s)}(x_1, \dots, x_s) \\ &= \sum_{n_1=0}^n \sum_{n_2=0}^{n-n_1} \dots \sum_{n_{s-1}=0}^{n-n_1-\dots-n_{s-2}} \mathbf{K}(n_1, \dots, n_{s-1}) F_{n-(n_1+\dots+n_{s-1}), n_1, \dots, n_{s-1}}^{(A_1, \dots, A_s; B_1, \dots, B_s)}(x_1, \dots, x_s) \\ & \quad \cdot \mathbf{N}(n_1, \dots, n_{s-1}); \end{aligned}$$

$$\begin{aligned} \mathbf{K}(n_1, \dots, n_{s-1}) &= \frac{(A_1+B_1+I)_{n-(n_1+\dots+n_{s-1})} (A_2+B_2+I)_{n_1} \dots (A_s+B_s+I)_{n_{s-1}}}{(c_1(a_1-b_1))^{n-(n_1+\dots+n_{s-1})} (c_2(a_2-b_2))^{n_1} \dots (c_s(a_s-b_s))^{n_{s-1}}}, \\ \mathbf{N}(n_1, \dots, n_{s-1}) &= [(A_s+I)_{n_{s-1}}]^{-1} [(A_{s-1}+I)_{n_{s-2}}]^{-1} \times \dots \\ & \quad \times [(A_2+I)_{n_1}]^{-1} [(A_1+I)_{n-(n_1+\dots+n_{s-1})}]^{-1} \end{aligned}$$

and also A_i and B_i are matrices in $\mathbb{C}^{r \times r}$ satisfying the spectral conditions $\text{Re}(z_i) > -1$ for each eigenvalue $z_i \in \sigma(A_i)$ and $\text{Re}(\eta_i) > -1$ for each eigenvalue $\eta_i \in \sigma(B_i)$, all matrices are commutative and $A_i + kI$ is invertible for every integer $k \geq 0$ for $1 \leq i \leq s$.

Theorem 6. For MEJMPs $F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x})$, we have hypergeometric matrix representation as follows:

$$F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) = \prod_{i=1}^s \left\{ \frac{(c_i(a_i-b_i))^{n_i}}{n_i!} F\left(A_i+B_i+(n_i+1)I, -n_i I; A_i+I; \frac{x_i-a_i}{b_i-a_i}\right) (A_i+I)_{n_i} \right\}.$$

In order to obtain some recurrence relations, we need the following lemma.

Lemma 1. Let a generating matrix function for $g_{n_1, \dots, n_s}(\mathbf{x}, \mathbf{C})$ be

$$(1-t_1)^{-C_1} \dots (1-t_s)^{-C_s} \Psi\left(\frac{-4x_1 t_1}{(1-t_1)^2}, \dots, \frac{-4x_s t_s}{(1-t_s)^2}\right) = \sum_{n_1, \dots, n_s=0}^{\infty} g_{n_1, \dots, n_s}(\mathbf{x}, \mathbf{C}) t_1^{n_1} \dots t_s^{n_s}$$

where $\mathbf{x} = (x_1, \dots, x_s)$, $\mathbf{C} = (C_1, \dots, C_s)$ and $g_{n_1, \dots, n_s}(\mathbf{x}, \mathbf{C})$ is a matrix polynomial of degree n_i with respect to x_i (of total degree $n = n_1 + \dots + n_s$), provided that

$$\begin{aligned} \Psi(u_1, \dots, u_s) &= \Psi_1(u_1) \dots \Psi_s(u_s); \quad u_i = \frac{-4x_i t_i}{(1-t_i)^2}, \quad i = 1, 2, \dots, s \\ \Psi_i(u_i) &= \sum_{n_i=0}^{\infty} \gamma_{n_i} u_i^{n_i}, \quad \gamma_0 \neq \mathbf{0} \end{aligned}$$

Then we have

$$\begin{aligned}
 & \text{i) } x_i \frac{\partial}{\partial x_i} g_{n_1, \dots, n_s}(\mathbf{x}, \mathbf{C}) - n_i g_{n_1, \dots, n_s}(\mathbf{x}, \mathbf{C}) \\
 &= -(C_i + (n_i - 1)I) g_{n_1, \dots, n_{i-1}, n_i - 1, n_{i+1}, \dots, n_s}(\mathbf{x}, \mathbf{C}) - x_i \frac{\partial}{\partial x_i} g_{n_1, \dots, n_{i-1}, n_i - 1, n_{i+1}, \dots, n_s}(\mathbf{x}, \mathbf{C}), \quad n_i \geq 1
 \end{aligned} \tag{11}$$

$$\begin{aligned}
 & \text{ii) } x_i \frac{\partial}{\partial x_i} g_{n_1, \dots, n_s}(\mathbf{x}, \mathbf{C}) - n_i g_{n_1, \dots, n_s}(\mathbf{x}, \mathbf{C}) \\
 &= -C_i \sum_{k=0}^{n_i-1} g_{n_1, \dots, n_{i-1}, k, n_{i+1}, \dots, n_s}(\mathbf{x}, \mathbf{C}) - 2x_i \sum_{k=0}^{n_i-1} \frac{\partial}{\partial x_i} g_{n_1, \dots, n_{i-1}, k, n_{i+1}, \dots, n_s}(\mathbf{x}, \mathbf{C}), \quad n_i \geq 1
 \end{aligned} \tag{12}$$

$$\begin{aligned}
 & \text{iii) } x_i \frac{\partial}{\partial x_i} g_{n_1, \dots, n_s}(\mathbf{x}, \mathbf{C}) - n_i g_{n_1, \dots, n_s}(\mathbf{x}, \mathbf{C}) = \\
 & \quad \sum_{k=0}^{n_i-1} (-1)^{n_i-k} (C_i + 2kI) g_{n_1, \dots, n_{i-1}, k, n_{i+1}, \dots, n_s}(\mathbf{x}, \mathbf{C}), \quad n_i \geq 1
 \end{aligned} \tag{13}$$

where $C_i C_j = C_j C_i$ for $i, j = 1, 2, \dots, s$.

Choosing

$$C_i = A_i + B_i + I \quad ; \quad \gamma_{n_i} = \frac{(I + A_i + B_i) 2^{n_i}}{2^{2n_i} n_i!} (I + A_i)_{n_i}^{-1} \tag{14}$$

in Lemma 1 and considering Theorem 2, we see that the matrix polynomials $g_{\mathbf{n}}$ is

$$\begin{aligned}
 g_{n_1, \dots, n_s}(\mathbf{v}) &= (\mathbf{c}(\mathbf{a} - \mathbf{b}))^{-\mathbf{n}} (\mathbf{A} + \mathbf{B} + I)_{\mathbf{n}} [(\mathbf{A} + I)_{\mathbf{n}}]^{-1} F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}((\mathbf{b} - \mathbf{a})\mathbf{v} + \mathbf{a}) \\
 &= \prod_{i=1}^s \left\{ (c_i (a_i - b_i))^{-n_i} (A_i + B_i + I)_{n_i} [(A_i + I)_{n_i}]^{-1} \right. \\
 & \quad \left. \cdot F_{n_i}^{(A_i, B_i)}((b_i - a_i)v_i + a_i; a_i, b_i, c_i) \right\}
 \end{aligned}$$

where A_i, B_i ($i = 1, 2, \dots, s$) are commutative matrices in $\mathbb{C}^{r \times r}$. With the help of Lemma 1 and also considering (11)-(13), one can easily obtain the next result.

Theorem 7. For the matrix polynomials $F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x})$, we have

$$\begin{aligned}
 & \text{i) } (x_i - a_i) \left[(A_i + B_i + n_i I) \frac{\partial F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x})}{\partial x_i} + c_i (a_i - b_i) \frac{\partial F_{n_1, \dots, n_{i-1}, n_i - 1, n_{i+1}, \dots, n_s}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x})}{\partial x_i} (A_i + n_i I) \right] \\
 &= (A_i + B_i + n_i I) \left[n_i F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) - c_i (a_i - b_i) F_{n_1, \dots, n_{i-1}, n_i - 1, n_{i+1}, \dots, n_s}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) (A_i + n_i I) \right], \\
 & \text{ii) } (x_i - a_i) \frac{\partial F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x})}{\partial x_i} - n_i F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) \\
 &= - \{c_i (a_i - b_i)\}^{n_i} (A_i + B_i + I)_{n_i}^{-1} \sum_{k=0}^{n_i-1} \left\{ (A_i + B_i + I)_k \{c_i (a_i - b_i)\}^{-k} \right. \\
 & \quad \left. \cdot \left\{ (A_i + B_i + I) F_{n_1, \dots, n_{i-1}, k, n_{i+1}, \dots, n_s}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) + 2(x_i - a_i) \frac{\partial F_{n_1, \dots, n_{i-1}, k, n_{i+1}, \dots, n_s}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x})}{\partial x_i} \right\} \right. \\
 & \quad \left. \cdot (I + A_i)_k^{-1} (A_i + I)_{n_i} \right\},
 \end{aligned}$$

$$\begin{aligned}
\text{iii) } (x_i - a_i) \frac{\partial F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x})}{\partial x_i} - n_i F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) &= \{c_i (b_i - a_i)\}^{n_i} (A_i + B_i + I)_{n_i}^{-1} \\
&\cdot \sum_{k=0}^{n_i-1} \left\{ \{c_i (b_i - a_i)\}^{-k} (A_i + B_i + (2k+1)I) (A_i + B_i + I)_k \right. \\
&\cdot \left. F_{n_1, \dots, n_{i-1}, k, n_{i+1}, \dots, n_s}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) (I + A_i)_k^{-1} (A_i + I)_{n_i} \right\},
\end{aligned}$$

where A_i and B_i are matrices in $\mathbb{C}^{r \times r}$ satisfying the spectral conditions $\text{Re}(z_i) > -1$ for every eigenvalue $z_i \in \sigma(A_i)$ and $\text{Re}(\eta_i) > -1$ for every eigenvalue $\eta_i \in \sigma(B_i)$, these matrices are commutative and $A_i + B_i + n_i I$ is invertible for every integer $n_i \geq 0$ for $1 \leq i \leq s$.

Remark 1. For the case of $s = 1$ in (9), extended Jacobi matrix polynomials (EJMPs) satisfy following equations [1]:

$$\begin{aligned}
(x-a) \left[(A+B+nI) \frac{d}{dx} F_n^{(A,B)}(x; a, b, c) + c(a-b) \frac{d}{dx} F_{n-1}^{(A,B)}(x; a, b, c) (A+nI) \right] \\
= (A+B+nI) \left[n F_n^{(A,B)}(x; a, b, c) - c(a-b) F_{n-1}^{(A,B)}(x; a, b, c) (A+nI) \right], \\
(x-a) \frac{d}{dx} F_n^{(A,B)}(x; a, b, c) - n F_n^{(A,B)}(x; a, b, c)
\end{aligned}$$

$$\begin{aligned}
&= -\{c(a-b)\}^n (A+B+I)_n^{-1} \sum_{k=0}^{n-1} \{c(a-b)\}^{-k} (A+B+I)_k \\
&\cdot \left\{ (A+B+I) F_k^{(A,B)}(x; a, b, c) + 2(x-a) \frac{d}{dx} F_k^{(A,B)}(x; a, b, c) \right\} \\
&\cdot (I+A)_k^{-1} (I+A)_n,
\end{aligned}$$

and

$$\begin{aligned}
(x-a) \frac{d}{dx} F_n^{(A,B)}(x; a, b, c) - n F_n^{(A,B)}(x; a, b, c) \\
= \{c(b-a)\}^n (A+B+I)_n^{-1} \sum_{k=0}^{n-1} \{c(b-a)\}^{-k} (A+B+(2k+1)I) \\
\cdot (A+B+I)_k F_k^{(A,B)}(x; a, b, c) (I+A)_k^{-1} (I+A)_n.
\end{aligned}$$

where all eigenvalues z of the matrices A and B of the extended Jacobi matrix polynomials $F_n^{(A,B)}(x; a, b, c)$ satisfy the condition $\text{Re}(z) > -1$ and $A+B+nI$ is invertible for every integer $n \geq 0$.

Remark 2. If we take $a = 1, b = -1$ and $c = -\frac{1}{2}$ in Theorem 7, we have recurrence relations for Jacobi matrix polynomials given in [30]. For the case $r = 1$, if we take $A_i = \alpha_i$ and $B_i = \beta_i$ with α_i, β_i real parameters and $\alpha_i, \beta_i > -1$ in Theorem 7, we have some recurrence relations satisfied by classical extended Jacobi polynomials $F_{\mathbf{n}}^{(\alpha_1, \dots, \alpha_s; \beta_1, \dots, \beta_s)}(\mathbf{x})$ given in [3].

4. MULTILINEAR AND MULTILATERAL GENERATING MATRIX FUNCTIONS

In recent years, by making use of the familiar group-theoretic (Lie algebraic) method a certain mixed trilateral finite-series relationships have been proved for orthogonal polynomials (see, for instance, [28]). In this section, we derive several

families of multilinear and multilateral generating matrix functions for the MEJMPs without using Lie algebraic techniques but, with the help of the similar method as considered in [1, 5].

Theorem 8. *Corresponding to a non-vanishing function $\Omega_\mu(y_1, \dots, y_r)$ consisting of r complex variables y_1, \dots, y_r ($r \in \mathbb{N}$) and of complex order μ , let*

$$\Lambda_{\mu,\nu}(y_1, \dots, y_r; z) := \sum_{k=0}^{\infty} a_k \Omega_{\mu+\nu k}(y_1, \dots, y_r) z^k \quad (15)$$

$(a_k \neq 0, \mu, \nu \in \mathbb{C})$

and

$$\begin{aligned} & \Theta_{\mathbf{n},p,\mu,\nu}(\mathbf{x}; y_1, \dots, y_r; \zeta) \\ & := \sum_{k=0}^{\lfloor n_1/p \rfloor} a_k \left\{ \prod_{i=2}^s (c_i (a_i - b_i))^{-n_i} \right\} (c_1 (b_1 - a_1))^{pk-n_1} F_{n_1-pk, n_2, \dots, n_s}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) \\ & \cdot \Omega_{\mu+\nu k}(y_1, \dots, y_r) \zeta^k \end{aligned} \quad (16)$$

where $\mathbf{n} = n_1 + \dots + n_s$; $n_1, \dots, n_s, p \in \mathbb{N}$, $\mathbf{x} = (x_1, \dots, x_s)$, $\mathbf{A} = (A_1, \dots, A_s)$, $\mathbf{B} = (B_1, \dots, B_s)$ and A_i and B_i are matrices in $\mathbb{C}^{r \times r}$ satisfying the spectral conditions $\text{Re}(z_i) > -1$ for every eigenvalue $z_i \in \sigma(A_i)$ and $\text{Re}(\eta_i) > -1$ for every eigenvalue $\eta_i \in \sigma(B_i)$ for $1 \leq i \leq s$ and all matrices are commutative. Then we have

$$\begin{aligned} & \sum_{n_1, \dots, n_s=0}^{\infty} \Theta_{\mathbf{n},p,\mu,\nu} \left(\mathbf{x}; y_1, \dots, y_r; \frac{\eta}{t_1^p} \right) t_1^{n_1} \dots t_s^{n_s} \\ & = \prod_{i=1}^s F_4 \left(I + B_i, I + A_i; I + A_i, I + B_i; \frac{(x_i - a_i) t_i}{a_i - b_i}, \frac{(x_i - b_i) t_i}{a_i - b_i} \right) \\ & \cdot \Lambda_{\mu,\nu}(y_1, \dots, y_r; \eta) \\ & \left(\sqrt{\frac{(x_i - a_i) t_i}{a_i - b_i}} + \sqrt{\frac{(x_i - b_i) t_i}{a_i - b_i}} < 1; i = 1, 2, \dots, s \right) \end{aligned} \quad (17)$$

provided that each member of (17) exists.

Proof. For convenience, let S denote the first member of the assertion (17) of Theorem 8. Then, plugging the polynomials

$$\Theta_{\mathbf{n},p,\mu,\nu} \left(\mathbf{x}; y_1, \dots, y_r; \frac{\eta}{t_1^p} \right)$$

from the definition (16) into the left-hand side of (17), we obtain

$$\begin{aligned} S & = \sum_{n_1, \dots, n_s=0}^{\infty} \sum_{k=0}^{\lfloor n_1/p \rfloor} a_k \left\{ \prod_{i=2}^s (c_i (a_i - b_i))^{-n_i} \right\} (c_1 (b_1 - a_1))^{pk-n_1} F_{n_1-pk, n_2, \dots, n_s}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) \\ & \cdot \Omega_{\mu+\nu k}(y_1, \dots, y_r) \eta^k t_1^{n_1-pk} t_2^{n_2} \dots t_s^{n_s}. \end{aligned} \quad (18)$$

Upon changing the order of summation in (18), if we replace n_1 by $n_1 + pk$, we can write

$$\begin{aligned}
S &= \sum_{n_1, \dots, n_s=0}^{\infty} \sum_{k=0}^{\infty} a_k \left\{ \prod_{i=1}^s (c_i (a_i - b_i))^{-n_i} \right\} F_{n_1, n_2, \dots, n_s}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) \Omega_{\mu+\nu k}(y_1, \dots, y_r) \eta^k t_1^{n_1} \dots t_s^{n_s} \\
&= \left(\sum_{n_1, \dots, n_s=0}^{\infty} \left\{ \prod_{i=1}^s (c_i (a_i - b_i))^{-n_i} \right\} F_{\mathbf{n}}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) t_1^{n_1} \dots t_s^{n_s} \right) \left(\sum_{k=0}^{\infty} a_k \Omega_{\mu+\nu k}(y_1, \dots, y_r) \eta^k \right) \\
&= \prod_{i=1}^s F_4 \left(I + B_i, I + A_i; I + A_i, I + B_i; \frac{(x_i - a_i) t_i}{a_i - b_i}, \frac{(x_i - b_i) t_i}{a_i - b_i} \right) \\
&\quad \cdot \Lambda_{\mu, \nu}(y_1, \dots, y_r; \eta)
\end{aligned}$$

which completes the proof of Theorem 8. \square

Theorem 9. *Corresponding to a non-vanishing function $\Omega_{\mu}(y_1, \dots, y_r)$ of r complex variables y_1, \dots, y_r ($r \in \mathbb{N}$) and of complex order μ , let*

$$\begin{aligned}
\Lambda_{\mu, \nu}(y_1, \dots, y_r; z) &:= \sum_{k=0}^{\infty} a_k \Omega_{\mu+\nu k}(y_1, \dots, y_r) z^k \\
&\quad (a_k \neq 0, \mu, \nu \in \mathbb{C})
\end{aligned} \tag{19}$$

and

$$\begin{aligned}
&\Theta_{n, p, \mu, \nu}(\mathbf{x}; y_1, \dots, y_r; \zeta) \\
&:= \sum_{k=0}^{\lfloor n/p \rfloor} a_k H_{n-pk}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) \Omega_{\mu+\nu k}(y_1, \dots, y_r) \zeta^k
\end{aligned} \tag{20}$$

where $n, p \in \mathbb{N}$, $\mathbf{x} = (x_1, \dots, x_s)$, $\mathbf{A} = (A_1, \dots, A_s)$, $\mathbf{B} = (B_1, \dots, B_s)$ and A_i and B_i are matrices in $\mathbb{C}^{r \times r}$ satisfying the spectral conditions $\operatorname{Re}(z_i) > -1$ for every eigenvalue $z_i \in \sigma(A_i)$ and $\operatorname{Re}(\eta_i) > -1$ for every eigenvalue $\eta_i \in \sigma(B_i)$ for $1 \leq i \leq s$. Then we have

$$\begin{aligned}
&\sum_{n=0}^{\infty} \Theta_{n, p, \mu, \nu} \left(\mathbf{x}; y_1, \dots, y_r; \frac{\eta}{t^p} \right) t^n \\
&= \prod_{i=1}^s (1-t)^{-(A_i + B_i + I)} F \left(\frac{A_i + B_i + I}{2}, \frac{A_i + B_i + 2I}{2}; A_i + I; \frac{4t(x_i - a_i)}{(a_i - b_i)(1-t)^2} \right) \\
&\quad \cdot \Lambda_{\mu, \nu}(y_1, \dots, y_r; \eta)
\end{aligned} \tag{21}$$

provided that each member of (21) exists. Here

$$\begin{aligned}
&H_k^{(A_1, \dots, A_s; B_1, \dots, B_s)}(x_1, \dots, x_s) \\
&= \sum_{k_1=0}^k \sum_{k_2=0}^{k-k_1} \dots \sum_{k_{s-1}=0}^{k-k_1-\dots-k_{s-2}} \mathbf{K}(k_1, \dots, k_{s-1}) F_{k-(k_1+\dots+k_{s-1}), k_1, \dots, k_{s-1}}^{(A_1, \dots, A_s; B_1, \dots, B_s)}(x_1, \dots, x_s) \mathbf{N}(k_1, \dots, k_{s-1});
\end{aligned}$$

and

$$\mathbf{K}(k_1, \dots, k_{s-1}) = \frac{(A_1 + B_1 + I)_{k-(k_1+\dots+k_{s-1})} (A_2 + B_2 + I)_{k_1} \dots (A_s + B_s + I)_{k_{s-1}}}{(c_1 (a_1 - b_1))^{k-(k_1+\dots+k_{s-1})} (c_2 (a_2 - b_2))^{k_1} \dots (c_s (a_s - b_s))^{k_{s-1}}},$$

$$\begin{aligned} \mathbf{N}(k_1, \dots, k_{s-1}) &= [(A_s + I)_{k_{s-1}}]^{-1} [(A_{s-1} + I)_{k_{s-2}}]^{-1} \times \dots \\ &\quad \times [(A_2 + I)_{k_1}]^{-1} [(A_1 + I)_{k-(k_1+\dots+k_{s-1})}]^{-1}, \end{aligned}$$

all matrices are commutative and $A_i + nI$ is invertible for every integer $n \geq 0$ for $1 \leq i \leq s$.

Proof. For convenience, let S denote the first member of the assertion (21) of Theorem 9. Then, upon substituting for the polynomials

$$\Theta_{n,p,\mu,\nu} \left(\mathbf{x}; y_1, \dots, y_r; \frac{\eta}{t^p} \right)$$

from the definition (20) into the left-hand side of (21), we obtain

$$S = \sum_{n=0}^{\infty} \sum_{k=0}^{[n/p]} a_k H_{n-pk}^{(A,B)}(\mathbf{x}) \Omega_{\mu+\nu k}(y_1, \dots, y_r) \eta^k t^{n-pk} \quad (22)$$

Upon changing the order of summation in (22), if we replace n by $n + pk$, we can write

$$\begin{aligned} S &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} a_k H_n^{(A,B)}(\mathbf{x}) \Omega_{\mu+\nu k}(y_1, \dots, y_r) \eta^k t^n \\ &= \left(\sum_{n=0}^{\infty} H_n^{(A,B)}(\mathbf{x}) t^n \right) \left(\sum_{k=0}^{\infty} a_k \Omega_{\mu+\nu k}(y_1, \dots, y_r) \eta^k \right) \\ &= \prod_{i=1}^s (1-t)^{-(A_i+B_i+I)} F \left(\frac{A_i+B_i+I}{2}, \frac{A_i+B_i+2I}{2}; A_i+I; \frac{4t(x_i-a_i)}{(a_i-b_i)(1-t)^2} \right) \\ &\quad \cdot \Lambda_{\mu,\nu}(y_1, \dots, y_r; \eta) \end{aligned}$$

which completes the proof of Theorem 9. \square

5. FURTHER CONSEQUENCES

By expressing the multivariable function $\Omega_{\mu+\nu k}(y_1, \dots, y_r)$, ($k \in \mathbb{N}_0$, $r \in \mathbb{N}$) in terms of a simpler function of one and more variables, we can give further applications of Theorem 8. For example, consider the case of $r = 1$ and $\Omega_{\mu+\nu k}(y) = L_{\mu+\nu k}^{(C,\lambda)}(y)$ in Theorem 8. Here, the Laguerre matrix polynomials $L_n^{(C,\lambda)}(y)$ are defined by [20] as follows:

$$L_n^{(C,\lambda)}(y) = \sum_{k=0}^n \frac{(-1)^k \lambda^k}{k! (n-k)!} (C+I)_n [(C+I)_k]^{-1} y^k$$

in which C is a matrix in $\mathbb{C}^{r \times r}$, $C + nI$ is invertible for every integer $n \geq 0$ and λ is a complex number with $Re(\lambda) > 0$. Notice that Laguerre matrix polynomials are generated by

$$\sum_{n=0}^{\infty} L_n^{(C,\lambda)}(y) t^n = (1-t)^{-(C+I)} \exp \left(\frac{-\lambda y t}{1-t} \right), \quad |t| < 1, \quad 0 < y < \infty. \quad (23)$$

Then we obtain the following result which provides a class of bilateral generating matrix functions for the MEJMPs and the Laguerre matrix polynomials.

Corollary 1. Let $\Lambda_{\mu,\nu}(y; z) := \sum_{k=0}^{\infty} a_k L_{\mu+\nu k}^{(C,\lambda)}(y) z^k$ where $(a_k \neq 0, \mu, \nu \in \mathbb{C})$ and

$$\begin{aligned} & \Theta_{\mathbf{n},p,\mu,\nu}(\mathbf{x}; y; \zeta) \\ & := \sum_{k=0}^{\lfloor n_1/p \rfloor} a_k \left\{ \prod_{i=2}^s (c_i (a_i - b_i))^{-n_i} \right\} (c_1 (b_1 - a_1))^{pk-n_1} F_{n_1-pk, n_2, \dots, n_s}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) L_{\mu+\nu k}^{(C,\lambda)}(y) \zeta^k \end{aligned}$$

where $\mathbf{x} = (x_1, \dots, x_s)$, $\mathbf{A} = (A_1, \dots, A_s)$, $\mathbf{B} = (B_1, \dots, B_s)$, $\mathbf{n} = n_1 + \dots + n_s$; $n_1, \dots, n_s, p \in \mathbb{N}$, $A_i B_i = B_i A_i$, $A_i A_j = A_j A_i$, $B_i B_j = B_j B_i$; $i, j = 1, 2, \dots, s$ and $C + nI$ is invertible for every integer $n \geq 0$. Then we have

$$\begin{aligned} & \sum_{n_1, \dots, n_s=0}^{\infty} \Theta_{\mathbf{n},p,\mu,\nu} \left(\mathbf{x}; y; \frac{\eta}{t_1^p} \right) t_1^{n_1} \dots t_s^{n_s} \\ & = \prod_{i=1}^s F_4 \left(I + B_i, I + A_i; I + A_i, I + B_i; \frac{(x_i - a_i) t_i}{a_i - b_i}, \frac{(x_i - b_i) t_i}{a_i - b_i} \right) \Lambda_{\mu,\nu}(y; \eta) \end{aligned} \quad (24)$$

provided that each member of (24) exists for $\sqrt{\frac{(x_i - a_i) t_i}{a_i - b_i}} + \sqrt{\frac{(x_i - b_i) t_i}{a_i - b_i}} < 1$; $i = 1, 2, \dots, s$.

Remark 3. Using the generating matrix function (23) for the Laguerre matrix polynomials and taking $a_k = 1$, $\mu = 0$, $\nu = 1$, we have

$$\begin{aligned} & \sum_{n_1, \dots, n_s=0}^{\infty} \sum_{k=0}^{\lfloor n_1/p \rfloor} \left\{ \prod_{i=2}^s (c_i (a_i - b_i))^{-n_i} \right\} (c_1 (b_1 - a_1))^{pk-n_1} F_{n_1-pk, n_2, \dots, n_s}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) \\ & \cdot L_k^{(C,\lambda)}(y) \eta^k t_1^{n_1-pk} t_2^{n_2} \dots t_s^{n_s} \\ & = \prod_{i=1}^s F_4 \left(I + B_i, I + A_i; I + A_i, I + B_i; \frac{(x_i - a_i) t_i}{a_i - b_i}, \frac{(x_i - b_i) t_i}{a_i - b_i} \right) \\ & \cdot (1 - \eta)^{-(C+I)} \exp \left(\frac{-\lambda y \eta}{1 - \eta} \right) \end{aligned}$$

where $|\eta| < 1$, $0 < y < \infty$ and $\sqrt{\frac{(x_i - a_i) t_i}{a_i - b_i}} + \sqrt{\frac{(x_i - b_i) t_i}{a_i - b_i}} < 1$ for $i = 1, \dots, s$.

Choose $r = 1$ and $\Omega_{\mu+\nu k}(y) = C_{\mu+\nu k}^D(y)$ where D is a matrix in $\mathbb{C}^{r \times r}$ satisfying the spectral condition

$$\left(\frac{-z}{2} \right) \notin \sigma(D) \text{ for } \forall z \in \mathbb{Z}^+ \cup \{0\}$$

in Theorem 8 where $\mu, \nu \in \mathbb{N}_0$. Here, Gegenbauer matrix polynomials are generated by [27] as follows:

$$\sum_{n=0}^{\infty} C_k^D(y) t^n = (1 - 2yt + t^2)^{-D} \quad , \quad |y| < 1 \quad (25)$$

Then we obtain a class of bilateral generating matrix functions for the MEJMPs and Gegenbauer matrix polynomials.

Corollary 2. Let $\Lambda_{\mu,\nu}(y; z) := \sum_{k=0}^{\infty} a_k C_{\mu+\nu k}^D(y) z^k$ where $(a_k \neq 0, \mu, \nu \in \mathbb{C})$ and

$$\begin{aligned} & \Theta_{\mathbf{n}, p, \mu, \nu}(\mathbf{x}; y; \zeta) \\ & := \sum_{k=0}^{\lfloor n_1/p \rfloor} a_k \left\{ \prod_{i=2}^s (c_i (a_i - b_i))^{-n_i} \right\} (c_1 (b_1 - a_1))^{pk-n_1} F_{n_1-pk, n_2, \dots, n_s}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) C_{\mu+\nu k}^D(y) \zeta^k \end{aligned}$$

where $\mathbf{x} = (x_1, \dots, x_s)$, $\mathbf{A} = (A_1, \dots, A_s)$, $\mathbf{B} = (B_1, \dots, B_s)$, $\mathbf{n} = n_1 + \dots + n_s$; $n_1, \dots, n_s, p \in \mathbb{N}$ and D is a matrix in $\mathbb{C}^{r \times r}$ satisfying the spectral condition

$$\left(\frac{-z}{2} \right) \notin \sigma(D) \text{ for } \forall z \in \mathbb{Z}^+ \cup \{0\},$$

A_i and B_i are matrices in $\mathbb{C}^{r \times r}$ satisfying the spectral conditions $\text{Re}(z_i) > -1$ for every eigenvalue $z_i \in \sigma(A_i)$ and $\text{Re}(\eta_i) > -1$ for every eigenvalue $\eta_i \in \sigma(B_i)$, $A_i B_i = B_i A_i$, $A_i A_j = A_j A_i$, $B_i B_j = B_j B_i$; $i, j = 1, 2, \dots, s$. Then we have

$$\begin{aligned} & \sum_{n_1, \dots, n_s=0}^{\infty} \Theta_{\mathbf{n}, p, \mu, \nu} \left(\mathbf{x}; y; \frac{\eta}{t_1^p} \right) t_1^{n_1} \dots t_s^{n_s} \\ & = \prod_{i=1}^s F_4 \left(I + B_i, I + A_i; I + A_i, I + B_i; \frac{(x_i - a_i) t_i}{a_i - b_i}, \frac{(x_i - b_i) t_i}{a_i - b_i} \right) \Lambda_{\mu, \nu}(y; \eta) \end{aligned} \quad (26)$$

provided that each member of (26) exists for $\sqrt{\frac{(x_i - a_i) t_i}{a_i - b_i}} + \sqrt{\frac{(x_i - b_i) t_i}{a_i - b_i}} < 1$; $i = 1, 2, \dots, s$.

Remark 4. Using (25) and taking $a_k = 1$, $\mu = 0$, $\nu = 1$, we have

$$\begin{aligned} & \sum_{n_1, \dots, n_s=0}^{\infty} \sum_{k=0}^{\lfloor n_1/p \rfloor} \left\{ \prod_{i=2}^s (c_i (a_i - b_i))^{-n_i} \right\} (c_1 (b_1 - a_1))^{pk-n_1} F_{n_1-pk, n_2, \dots, n_s}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) \\ & \cdot C_k^D(y) \eta^k t_1^{n_1-pk} t_2^{n_2} \dots t_s^{n_s} \\ & = \prod_{i=1}^s F_4 \left(I + B_i, I + A_i; I + A_i, I + B_i; \frac{(x_i - a_i) t_i}{a_i - b_i}, \frac{(x_i - b_i) t_i}{a_i - b_i} \right) \\ & \cdot (1 - 2y\eta + \eta^2)^{-D}. \end{aligned}$$

Setting $r = s$ and $\Omega_{\mu+\nu k}(y_1, \dots, y_s) = H_{\mu+\nu k}^{(C, D)}(y_1, \dots, y_s)$ in Theorem 9, we obtain result which provides a class of bilinear generating matrix functions for the MEJMPs.

Corollary 3. Let $\Lambda_{\mu,\nu}(y_1, \dots, y_s; z) := \sum_{k=0}^{\infty} a_k H_{\mu+\nu k}^{(C, D)}(\mathbf{y}) z^k$ where $(a_k \neq 0, \mu, \nu \in \mathbb{C})$ and

$$\Theta_{\mathbf{n}, p, \mu, \nu}(\mathbf{x}; \mathbf{y}; \zeta) := \sum_{k=0}^{\lfloor n/p \rfloor} a_k H_{n-pk}^{(\mathbf{A}, \mathbf{B})}(\mathbf{x}) H_{\mu+\nu k}^{(C, D)}(\mathbf{y}) \zeta^k \quad (27)$$

where $n, p \in \mathbb{N}$, $\mathbf{x} = (x_1, \dots, x_s)$, $\mathbf{y} = (y_1, \dots, y_s)$, $\mathbf{A} = (A_1, \dots, A_s)$, $\mathbf{B} = (B_1, \dots, B_s)$, $\mathbf{C} = (C_1, \dots, C_s)$, $\mathbf{D} = (D_1, \dots, D_s)$ and A_i, B_i, C_i and D_i ($i = 1, 2, \dots, s$) are matrices in $\mathbb{C}^{r \times r}$ whose eigenvalues, z , all satisfy $\text{Re}(z) > -1$. Then we have

$$\begin{aligned} & \sum_{n=0}^{\infty} \Theta_{n,p,\mu,\nu} \left(\mathbf{x}; \mathbf{y}; \frac{\eta}{t^p} \right) t^n \\ &= \prod_{i=1}^s (1-t)^{-(A_i+B_i+I)} F \left(\frac{A_i+B_i+I}{2}, \frac{A_i+B_i+2I}{2}; A_i+I; \frac{4t(x_i-a_i)}{(a_i-b_i)(1-t)^2} \right) \\ & \quad \cdot \Lambda_{\mu,\nu}(y_1, \dots, y_s; \eta) \end{aligned} \quad (28)$$

provided that each member of (28) exists.

Remark 5. Taking $a_k = 1$, $\mu = 0$, $\nu = 1$ and using Theorem 5, we have

$$\begin{aligned} & \sum_{n=0}^{\infty} \sum_{k=0}^{[n/p]} H_{n-pk}^{(\mathbf{A},\mathbf{B})}(\mathbf{x}) H_k^{(\mathbf{C},\mathbf{D})}(\mathbf{y}) \eta^k t^{n-pk} \\ &= \prod_{i=1}^s (1-t)^{-(A_i+B_i+I)} F \left(\frac{A_i+B_i+I}{2}, \frac{A_i+B_i+2I}{2}; A_i+I; \frac{4t(x_i-a_i)}{(a_i-b_i)(1-t)^2} \right) \\ & \quad \cdot \prod_{i=1}^s (1-\eta)^{-(C_i+D_i+I)} F \left(\frac{C_i+D_i+I}{2}, \frac{C_i+D_i+2I}{2}; C_i+I; \frac{4\eta(y_i-a_i)}{(a_i-b_i)(1-\eta)^2} \right) \end{aligned}$$

where

$$\begin{aligned} & H_k^{(C_1, \dots, C_s; D_1, \dots, D_s)}(y_1, \dots, y_s) \\ &= \sum_{k_1=0}^k \sum_{k_2=0}^{k-k_1} \dots \sum_{k_{s-1}=0}^{k-k_1-\dots-k_{s-2}} \mathbf{K}(k_1, \dots, k_{s-1}) F_{k-(k_1+\dots+k_{s-1}), k_1, \dots, k_{s-1}}^{(C_1, \dots, C_s; D_1, \dots, D_s)}(y_1, \dots, y_s) \mathbf{N}(k_1, \dots, k_{s-1}); \end{aligned}$$

and

$$\mathbf{K}(k_1, \dots, k_{s-1}) = \frac{(C_1 + D_1 + I)_{k-(k_1+\dots+k_{s-1})} (C_2 + D_2 + I)_{k_1} \dots (C_s + D_s + I)_{k_{s-1}}}{(c_1(a_1 - b_1))^{k-(k_1+\dots+k_{s-1})} (c_2(a_2 - b_2))^{k_1} \dots (c_s(a_s - b_s))^{k_{s-1}}},$$

$$\begin{aligned} \mathbf{N}(k_1, \dots, k_{s-1}) &= [(C_s + I)_{k_{s-1}}]^{-1} [(C_{s-1} + I)_{k_{s-2}}]^{-1} \times \dots \\ & \quad \times [(C_2 + I)_{k_1}]^{-1} [(C_1 + I)_{k-(k_1+\dots+k_{s-1})}]^{-1} \end{aligned}$$

and $C_i D_i = D_i C_i$, $C_i C_j = C_j C_i$, $D_i D_j = D_j D_i$; $i, j = 1, 2, \dots, s$ and $C_i + nI$ is invertible for every integer $n \geq 0$ for $1 \leq i \leq s$.

Furthermore, for every suitable choice of the coefficients a_k ($k \in \mathbb{N}_0$), if the multivariable function $\Omega_{\mu+\nu k}(y_1, \dots, y_r)$, ($r \in \mathbb{N}$), is expressed as an appropriate product of several simpler functions, the assertions of Theorem 8 and Theorem 9 can be applied in order to derive various families of multilinear and multilateral generating matrix functions for the MEJMPs.

REFERENCES

- [1] R. Aktaş, B. Çekim, A. Çevik, *Extended Jacobi matrix polynomials*, Util. Math. (in press).
- [2] R. Aktaş, B. Çekim, R. Şahin, *The matrix version for the multivariable Humbert polynomials*, Miskolc Mathematical Notes **13** (2) (2012), 197-208.
- [3] A. Altın, R. Aktaş, E. Erkuş-Duman, *On a multivariable extension for the extended Jacobi polynomials*, J. Math. Anal. Appl. **353** (2009), 121-133.

- [4] A. Altın, B. Çekim, *Some properties associated with Hermite matrix polynomials*, Util. Math. **88** (2012), 171-181.
- [5] A. Altın, B. Çekim, E. Erkuş-Duman, *Families of generating functions for the Jacobi and related matrix polynomials*, Ars Combin. (*in press*).
- [6] R.S. Batahan, *A new extension of Hermite matrix polynomials and its applications*, Linear Algebra Appl. **419** (2006) 82-92.
- [7] M.M. Castro, F.A. Grünbaum, *The Algebra of Differential Operators Associated to a Family of Matrix-Valued Polynomials: Five Instructive Examples*, IMRN, (2006), 1-33.
- [8] R.D. Costin, *Matrix valued polynomials generated by the scalar-type Rodrigues formulas*, J. Approx. Theory **162** (1) (2010), 141-152.
- [9] B. Çekim, A. Altın, R. Aktaş, *Some relations satisfied by orthogonal matrix polynomials*, Hacettepe Journal of Mathematics and Statistics **40** (2) (2011), 241-253.
- [10] E. Defez, L. Jódar, *Some applications of the Hermite matrix polynomials series expansions*, J. Comp. Appl. Math. **99** (1998), 105-117.
- [11] E. Defez, L. Jódar, *Chebyshev matrix polynomials and second order matrix differential equations*, Util. Math. **61** (2002), 107-123.
- [12] E. Defez, L. Jódar, A. Law, *Jacobi matrix differential equation, polynomial solutions and their properties*, Comput. Math. Appl. **48** (2004), 789-803.
- [13] N. Dunford, J. Schwartz, *Linear Operators. Vol. I*, Interscience, New York, 1957.
- [14] C.F. Dunkl, Y. Xu, *Orthogonal Polynomials in Several Variables*, Encyclopedia of Mathematics and its Applications, 81, Cambridge University Press, Cambridge 2001.
- [15] A.J. Durán, F.A. Grünbaum, *Orthogonal matrix polynomials, scalar-type Rodrigues formulas and Pearson equations*, J. Approx. Theory **134** (2005), 267-280.
- [16] I. Gohberg, P. Lancaster, L. Rodman, *Matrix Polynomials*, Academic Press, New York, 1982.
- [17] F.A. Grünbaum, *Matrix valued Jacobi polynomials*, Bulletin des Sciences Mathématiques **127** (3) (2003), 207-214.
- [18] F.A. Grünbaum, I. Pacharoni, J.A. Tirao, *Matrix valued orthogonal polynomials of the Jacobi type*, Indag. Math. (N.S.) **14** (3-4) (2003), 353-366.
- [19] L. Jódar, R. Company, *Hermite matrix polynomials and second order matrix differential equations*, J. Approx. Theory Appl. **12** (2) (1996), 20-30.
- [20] L. Jódar, R. Company, E. Navarro, *Laguerre matrix polynomials and system of second-order differential equations*, Appl. Num. Math. **15** (1994), 53-63.
- [21] L. Jódar, R. Company, E. Ponsoda, *Orthogonal matrix polynomials and systems of second order differential equations*, Differential Equations and Dynamical Systems **3** (1996), 269-288.
- [22] L. Jódar and J.C. Cortés, *On the hypergeometric matrix function*, J. Comp. Appl. Math. **99** (1998), 205-217.
- [23] L. Jódar, J.C. Cortés, *Closed form general solution of the hypergeometric matrix differential equation*, Math. Comput. Modelling **32** (2000), 1017-1028.
- [24] T.H. Koornwinder, *Two variable analogues of the classical orthogonal polynomials*. Theory and Application of Special Functions, Acad. Press. Inc., New York, 1975.
- [25] M.G. Krein, *Fundamental aspects of the representation theory of hermitian operators with deficiency index (m,m)* , AMS Translations, Series **2**, **97**, Providence, Rhode Island (1971), 75-143.
- [26] M.G. Krein, *Infinite J -matrices and a matrix moment problem*, Dokl. Akad. Nauk SSSR **2** (1949), 125-128.
- [27] K.A.M. Sayyed, M.S. Metwally, R.S. Batahan, *Gegenbauer matrix polynomials and second order matrix differential equations*, Divulg. Mat. **12** (2004), 101-115.
- [28] H.M. Srivastava, H.L. Manocha, *A Treatise on Generating Functions*, Halsted Press, John Wiley and Sons, New York, 1984.
- [29] P.K. Suetin, *Orthogonal Polynomials in Two Variables*, Gordon and Breach, Amsterdam, 1999.
- [30] F. Taşdelen, B. Çekim, R. Aktaş, *On a multivariable extension of Jacobi matrix polynomials*, Comput. Math. Appl. **61**(9) (2011), 2412-2423.

ALÍ ÇEVİK, MERSIN UNIVERSITY, DEPARTMENT OF MATHEMATICS, ÇİFTLİKKÖY TR-33343
MERSIN, TURKEY

E-mail address: cevik@mersin.edu.tr