

Relationship Between Seminal Plasma Composition and Spermatological Parameters in Scaly Carp (*Cyprinus carpio*)

¹Yusuf Bozkurt, ²Fatih Ogretmen, ³Faik Sertel Secer and ⁴Ugur Ercin

¹Department of Aquaculture, Faculty of Fisheries, University of Mustafa Kemal, Hatay, Turkey

²State Hydrolic Works, Fish Production Station, Adana, Turkey

³Department of Fisheries and Aquaculture, Faculty of Agriculture, University of Ankara, Ankara, Turkey

⁴Department of Biochemistry, Faculty of Medicine, University of Gazi, Ankara, Turkey

Abstract: The present study was conducted to evaluate seminal plasma composition, spermatological parameters and their relationships in scaly carp (*Cyprinus carpio*). Seminal plasma contained 71.18 ± 3.40 mML⁻¹ (Na⁺), 87.48 ± 4.42 mML⁻¹ (K⁺), 10.64 ± 1.50 mg dL⁻¹ (Ca²⁺), 2.57 ± 0.19 mEq L⁻¹ (Mg²⁺), 0.12 ± 0.04 g dL⁻¹ total protein, 1.43 ± 0.02 mg dL⁻¹ glucose, 7.82 ± 2.60 mg dL⁻¹ triglyceride, 6.27 ± 2.32 mg dL⁻¹ cholesterol, 24.45 ± 7.98 mg dL⁻¹ urea. The following spermatological parameters were found: sperm volume 13.17 ± 5.39 mL, sperm motility $63.18 \pm 7.16\%$, movement duration 56.81 ± 20.32 s, sperm concentration $29.637 \pm 1.07 \times 10^9$ mL⁻¹, total sperm concentration $419.200 \pm 29.62 \times 10^9$ and pH 7.59 ± 0.74 . The K⁺ and Ca²⁺ ions correlated positively with motility ($r = 0.215$, $p > 0.05$ and $r = 0.048$, $p > 0.05$), respectively. On the other hand, Na⁺ and Mg²⁺ ions correlated negatively with motility ($r = -0.067$, $p > 0.05$, $r = -0.407$, $p > 0.05$), respectively. Significant correlations were observed between K⁺ and Na⁺ ($r = -0.695$, $p < 0.05$), protein and triglyceride ($r = 0.786$, $p < 0.01$), protein and cholesterol ($r = 0.936$, $p < 0.01$), triglyceride and cholesterol ($r = 0.802$, $p < 0.01$). These parameters can be used to prepare species-specific extenders for short-term storage and cryopreservation of scaly carp sperm.

Key words: Scaly carp, semen, seminal plasma, sperm quality

INTRODUCTION

Broodfish management is an important factor that influence gamete quality and success of artificial insemination (Billard *et al.*, 1995). Better knowledge of sperm biochemistry, physiology and the mechanisms of reproduction are essential to improve artificial fertilization procedures for fish.

Evaluation of sperm quality provides information allowing aquaculturists to determine the optimal time for sperm collection and to devise optimal handling and storage protocols for sperm used in artificial fertilization (Billard *et al.*, 1995; Linhart *et al.*, 2004). Sperm volume, concentration and motility, as well as composition of the seminal plasma are common parameters to assess sperm quality in fish (Alavi and Cosson, 2006; Linhart *et al.*, 2000).

The composition of seminal plasma has great influence on the biological quality of the milt, as expressed by sperm viability and motility. These factors are directly related to the fertilization success. Determining of seminal plasma composition can help to

understand the design requirements for preparing the appropriate artificial seminal plasma solutions for short-term storage or cryopreservation.

In this regard, the inorganic compounds provide isotonic conditions for spermatozoa and especially, K⁺ ions prevent sperm motility in some freshwater fish species such as salmonids and sturgeons (Morisawa *et al.*, 1983). Also, some organic compounds including glucose and triglyceride act as the main energy resources for spermatozoa metabolism (Lahnsteiner *et al.*, 1993) and some others such as proteins prolong the viability of spermatozoa (Lahnsteiner *et al.*, 2004). To increase the efficiency of artificial fertilization, the composition of sperm diluents is very important and must be according to the species-specific composition of the seminal plasma.

Compared to the physiology of reproduction of other fishes, less attention has been paid to that of the scaly carp. Based on the literature, sperm characteristics of scaly carp has been rarely investigated. So, this study was designed to determine the physical and biochemical characteristics of scaly carp sperm and to investigate their physiological relationships with each other.

MATERIALS AND METHODS

Broodstock care and collection of semen: The experiment was carried out at the State Hydrolic Works (SHW) Fish Reproduction Station. The male broodstock were held in sand ponds (water temperature: 23-24°C, O₂: 7-8 ppm) under a natural photoperiod regime and fasted 48 h before semen collection. Eleven mature scaly carp males (TW = 2.45±1.24 kg, TL = 46.23±2.52 cm) were randomly selected from the stock and were used as semen donors in the middle of the spawning season. For sperm collection, fish were not fed 48 h prior to collection and the males were anaesthetized in 5 mg L⁻¹ of quinaldine sulphate before stripping. Before collection of sperm, the genital area was cleaned with a towel and dried to avoid contamination of sperm with feces, urine and water. Then, a gentle abdominal pressure was applied 12 h after a single injection of 2 mg⁻¹ body weight of Carp Pituitary Extract (CPE) at 23-24°C water temperature. Collected sperm was immediately transported to the laboratory for analyses.

Evaluation of semen: Sperm was sampled into the calibrated glass tubes and the volume was expressed as mL. Sperm motility was estimated from freshly collected samples of milt according to the percentage of motile spermatozoa. For this aim, a 5 µL drop of sperm was placed directly on a microscope slide and 100 µL of activation solution (0.3% NaCl) was added. Microscopic observation at 400x magnification was carried out at room temperature (20°C). Three fields of view were examined for each slide and three aliquots of each milt sample were inspected for calculation of an average. Duration of spermatozoa movement was assessed using a sensitive chronometer (1/100) that was started simultaneously with the addition of activation solution into the samples.

Spermatozoa concentration was determined according to the haemocytometric method. For this aim, sperm was diluted at ratio of 1:1000 with Hayem solution (5 g Na₂SO₄, 1 g NaCl, 0.5g HgCl₂, 200 mL bidistilled water). Ten microliter of the dilution were taken for counting on a Thoma hemocytometer slide (depth 0.1 mm). To prevent the spermatozoa adhering to the slides, they were previously bathed in a 10% bovine serum albumin solution (30 min, 4°C), then washed with distilled water and dried in a hood. Mean spermatozoa count was calculated from three replicate samples for each fish at magnification of 400x and expressed as ×10⁹ mL⁻¹. Sperm pH was measured using standard pH papers (Merck) within 30 min of sampling. Semen colour was evaluated visually immediately following sperm collection.

Determination of seminal plasma composition: Seminal plasma was collected after centrifugation of semen at 4000× g for 10 min at room temperature (20°C) and stored in Eppendorf (Wiesbaden, Germany) vials at -20°C, until the beginning of the analyses. Seminal plasma was centrifuged twice to avoid possible contamination with spermatozoa. Major cations and metabolites such as Na⁺, K⁺, Ca²⁺, Mg²⁺, Cl⁻, glucose, protein, cholesterol, triglyceride and urea levels were determined using an Abbott-Aeroset autoanalyser (Chicago, USA) using original kits.

Statistical analysis: Motility data were normalized through arcsine transformation. Correlations between physical spermatological parameters and seminal plasma composition were estimated using Pearson's correlation test. Results are presented as mean±SEM. Statistical analyses were performed with SPSS 10 for Windows statistical software package.

RESULTS

Spermatological parameters: Spermatological parameters of the collected sperm were found to be rather variable and are presented in Table 1. The sperm volume collected for each male ranged between 7.4 and 23.4 mL and the mean was found to be 13.17±5.39 mL. Sperm volume correlated significantly with motility (r = 0.653, p<0.05) and total concentration (r = 0.834, p<0.01). The spermatozoa motility ranged between 60 and 75% and the mean was found to be 63.18±7.16. Sperm was found to be viscous in consistency and creamy white in colour in all samples.

Seminal plasma composition: The seminal plasma ionic and metabolite composition are shown in Table 2. Main

Table 1: Spermatological parameters of scaly carp (n = 11)

Parameters	Minimum	Maximum	Mean	SEM
Volume (mL)	7.400	23.400	13.170	5.39
Motility (%)	60.000	75.000	63.180	7.16
Movement duration (s)	28.000	85.000	56.810	20.32
Density (×10 ⁹ mL ⁻¹)	20.270	52.750	29.637	1.07
Total density (×10 ⁹)	200.100	967.200	419.200	29.62
pH	6.900	9.200	7.590	0.74

Table 2: Seminal plasma ion and metabolite composition of scaly carp sperm (n = 3)

Variables	Minimum	Maximum	Mean	SEM
Na ⁺ (mM L ⁻¹)	68.1	77.4	71.18	3.40
K ⁺ (mM L ⁻¹)	81.4	97.1	87.48	4.42
Ca ²⁺ (mg dL ⁻¹)	7.5	13.3	10.64	1.50
Mg ²⁺ (mEq L ⁻¹)	2.3	2.9	2.57	0.19
Glucose (mg dL ⁻¹)	1.2	1.8	1.43	0.02
Total Protein (g dL ⁻¹)	0.1	0.4	0.12	0.04
Triglyceride (mg dL ⁻¹)	5.0	12.0	7.82	2.60
Cholesterol (mg dL ⁻¹)	5.0	11.0	6.27	2.32
Urea (mg dL ⁻¹)	10.0	37.0	24.45	7.98

Table 3: Correlations between spermatological parameters and seminal plasma composition of scaly carp sperm

Variables	Volume	Movement		Total		pH	Ca	Na	K	Mg	Glucose	Protein	Triglyceride	Cholesterol
		Motility	duration	Density	density									
Motility	0.653*													
Movement duration	-0.444	0.542												
Density	0.549	0.417	-0.293											
Total density	0.834**	0.602	-0.408	0.914**										
pH	-0.219	-0.013	0.359	0.049	-0.123									
Ca	-0.105	0.048	0.277	-0.199	-0.152	0.054								
Na	0.330	-0.067	0.145	-0.282	-0.043	-0.127	-0.249							
K	-0.256	0.215	-0.559	-0.149	-0.209	-0.112	0.147	-0.695*						
Mg	0.013	-0.407	0.198	-0.496	-0.304	-0.417	0.213	0.271	-0.164					
Glucose	0.165	0.376	-0.199	-0.069	0.052	-0.040	0.396	0.061	0.069	0.137				
Protein	-0.001	-0.285	-0.353	-0.274	-0.202	0.008	-0.015	0.155	0.129	0.430	0.261			
Triglyceride	-0.236	-0.234	-0.467	-0.360	-0.361	-0.079	-0.024	0.027	0.418	0.029	0.141	0.786**		
Cholesterol	-0.269	-0.537	-0.155	-0.350	-0.379	0.060	-0.089	0.107	0.072	0.380	0.052	0.936**	0.802**	
Urea	0.183	-0.028	-0.447	-0.157	-0.047	-0.001	-0.545	0.162	0.186	0.273	-0.343	0.527	0.318	0.472

*Significant at $p < 0.05$, **Significant at $p < 0.01$

ionic components such as Na^+ , K^+ , Ca^{2+} and Mg^{2+} were found rather variable. There were no significant difference among Ca^{2+} , K^+ and motility ($r = 0.048$, $p > 0.05$, $r = 0.215$, $p > 0.05$) respectively. On the other hand, negative relationship was determined between motility, Na^+ and Mg^{2+} ions ($r = -0.067$, $p > 0.05$; $r = -0.407$, $p > 0.05$) (Table 3).

Spermatozoa motility showed negative relationship with the metabolites of the seminal plasma such as protein, triglyceride, cholesterol and urea ($r = -0.285$, $p > 0.05$; $r = -0.234$, $p > 0.05$; $r = -0.537$, $p > 0.05$; $r = -0.028$, $p > 0.05$), but positive relationship was determined with glucose ($r = 0.376$, $p > 0.05$), respectively. On the other hand, spermatozoa concentration showed negative allometry with all ionic and metabolic composition of the seminal plasma ($p > 0.05$). Protein significantly correlated with triglyceride and cholesterol ($r = 0.786$, $p < 0.01$; $r = 0.936$, $p < 0.01$). A highly significant relationship was also found between triglyceride and cholesterol ($r = 0.802$, $p < 0.01$).

DISCUSSION

Mean semen volume was similar with the findings of Akcay *et al.* (2004) for mirror carp but differed from results reported by Bozkurt and Secer (2005) for mirror carp. The differences may be due to the feeding conditions and regime, water quality, environmental factors or spawning time. In the present study, mean sperm motility and motility duration was determined as $63.18 \pm 7.16\%$ and 56.81 ± 20.32 sec, respectively. Spermatozoa motility varies in vigor and duration not only among males but also within an individual male depending on ripeness (Tekin *et al.*, 2003) and probably due to contamination with urine during stripping. Also, most studies on fish species have shown that the duration and motility of semen can vary seasonally (Akcay *et al.*, 2004; Benau and Turner, 1980). Spermatozoa concentration showed great differences with the findings of Emri *et al.* (1998),

Akcay *et al.* (2004) and Bozkurt and Secer (2005). The differences may be due to differences in feeding conditions, age, environmental factors, time of spawning, or dilution ratio.

According to the results of the present study, seminal plasma of scaly carp has lower Na^+ content (71.18 ± 3.40 mM L^{-1}) than other freshwater species such as grass carp (98 mmol L^{-1}) (Bozkurt *et al.*, 2008), rainbow trout (80 mmol L^{-1}) (Secer *et al.*, 2004), perch (124 mmol L^{-1}) (Lahnsteiner *et al.*, 1995). On the other hand, the concentration of K^+ (87.48 ± 4.42 mM L^{-1}) in the seminal plasma was higher than in Atlantic salmon (28 mM L^{-1}) (Aas *et al.*, 1991), *Salmo trutta abanticus* (38 mM L^{-1}) (Bozkurt, 2008), rainbow trout (46 mM L^{-1}) (Secer *et al.*, 2004) and perch (10 mM L^{-1}) (Lahnsteiner *et al.*, 1995).

The electrolytes ensure the viability of sperm. Low level of the Na^+ ion can be associated with low percentages of motility and may be caused by a deficiency in the formation of seminal plasma. The K^+ ion has a specific role in maintaining spermatozoa in the quiescent state (Baynes *et al.*, 1981). K^+ has an inhibitory effect on the initiation of sperm motility in salmonids (Billard and Cosson, 1992). Similarly, findings of the present study agree with the inhibitory effect of K^+ ion on motility like in salmonids. The percentage of motile cells of scaly carp spermatozoa are observed tend to increase when, the Mg^{2+} ion level in decrease. It seems that Mg^{2+} at certain concentrations tends to inhibit the spermatozoa in scaly carp.

The specific role of protein in fish semen is unknown. White and Macleod (1963) indicated that protein had a protective role. In this study, low concentrations of total protein (0.12 ± 0.04 g dL^{-1}) were found, which indicates a low demand for protein. However, because of the negative relationship of protein with the sperm motility the possible role of protein remains underfined. On the other hand, Lahnsteiner *et al.* (2004) found that seminal plasma proteins prolong the viability of rainbow trout

spermatozoa as measured by sperm motility. Notable concentrations of urea (24.45 ± 7.98 mg dL⁻¹) were also found in the seminal plasma. Urea is considered in relationship with protein metabolism and total protein.

Fish spermatozoa are capable of utilizing extracellular carbohydrates. In this study, glucose has been identified in the seminal plasma and its concentration was found to be 1.43 ± 0.02 mg dL⁻¹. The importance of glucose in fish semen is not clear. On the other hand, the presence of this sugar in seminal plasma has been connected to the high energy demand of the testes during spermatogenesis or for the lipid synthesis of spermatozoa (Soengas *et al.*, 1993).

Various lipid classes have been found in seminal plasma and their levels are highly variable among fish species, such as 0.007 g L⁻¹ for Arctic charr and 1.00 g L⁻¹ for Euroasian perch (Piironen and Hyvarinen, 1983). In the present study, the mean level of triglyceride (7.82 ± 2.60 mg dL⁻¹) was negatively correlated with sperm motility. Triglycerides serve as energy resources for spermatozoa during immotile storage and during the regeneration phase after motility (Lahnsteiner *et al.*, 1993). According to Lahnsteiner *et al.* (1994), low levels of triglycerides were found in the seminal plasma of cyprinids. Low triglyceride levels could therefore be indicative of inadequate energy resources, reduced motility rate and fertilization capacity.

Also, the cholesterol level was found to be 6.27 ± 2.32 mg dL⁻¹ in this study. In spite of the identification of cholesterol in the seminal plasma of freshwater fish (Billard *et al.*, 1995), there is not enough information about its role. Lipids and cholesterol probably might have a protective effect on environmental changes (especially on temperature) that might occur when fish semen are released.

CONCLUSION

The results of this study confirmed differences in the compositions of seminal plasma in related species. The greatest changes in the ionic composition of the seminal plasma in scaly carp has been noted in the K⁺ ion concentrations. This study also, suggests that the concentration of the organic and inorganic compounds in artificial seminal plasma should be modified to meet species specific extender requirements for short-term storage or cryopreservation of scaly carp sperm.

ACKNOWLEDGMENTS

The researchers are grateful to the staff of the SHW Fish Production Station, Adana, Turkey. The researchers also, would like to thank Dr. N. Bukan for providing necessary materials and conditions for seminal plasma analysis.

REFERENCES

- Aas, G.H., T. Refstie and B. Gjerde, 1991. Evaluation of milt quality of Atlantic salmon. *Aquaculture*, 95: 125-132. DOI: 10.1016/0044-8486(91)90079-M.
- Akcay, E., Y. Bozkurt, S. Secer, N. Tekin, 2004. Cryopreservation of mirror carp semen. *Turk. J. Vet. Anim. Sci.*, 28: 837-843. <http://journals.tubitak.gov.tr/veterinary/issues/vet-04-28-5/vet-28-5-9-0211-37.pdf>.
- Alavi, S.M.H. and J. Cosson, 2006. Sperm motility in fishes: (II) effects of ions and osmotic pressure. *Cell. Biol. Int.*, 30: 1-14. DOI: 10.1016/j.cellbi.2005.06.004.
- Baynes, S.M., A.P. Scott and A.P. Dawson, 1981. Rainbow trout, *Salmo gairdneri* Richardson, spermatozoa: Effects of cations and pH on motility. *J. Fish Biol.*, 19: 259-267. DOI: 10.1111/j.1095-8649.1981.tb05830.
- Benau, D. and C. Termer, 1980. Initiation, prolongation and reactivation of the motility of salmonid spermatozoa. *Gamete Res.*, 3 (3): 247-257. OCLC: 3531657.
- Billard, R. and M.P. Cosson, 1992. Some problems related to the assessment of sperm motility in freshwater fish. *J. Exp. Zool.*, 261: 122-131. <http://www3.interscience.wiley.com/search/allsearch?mode=citation&contextLink=blah&issn=%281932-5231%2C1552-499X%2C1552499X%29&volume=261&issue=&pages=122>.
- Billard, R., J. Cosson, L.W. Crim and M. Suquet, 1995. Sperm Physiology and Quality. In: Bromage, N.R. and R.J. Roberts (Eds.). *Brood Stock Management and Egg and Larval Quality*, Blackwell Science, Oxford, UK, pp: 25-52. ISBN: 0-632-03591-9.
- Bozkurt, Y. and S. Secer, 2005. Effect of short term preservation of mirror carp (*Cyprinus carpio*) semen on motility, fertilization and hatching rates. *Isr. J. Aquac. Bamidgeh*, 57 (3): 125-130. <http://www.siamb.org.il/uploads/573bozkurt.abs.pdf>.
- Bozkurt, Y., 2008. Physical and biochemical properties of *Salmo trutta abanticus* semen. *Indian Vet. J.*, 85 (3): 282-284. IDS Number: 295XH.
- Bozkurt, Y., F. Oğretmen, U. Ercin and U. Yildiz, 2008. Seminal plasma composition and its relationship with physical spermatological parameters of grass carp (*Ctenopharyngodon idella*) semen: With emphasis on sperm motility. *Aquac. Res.*, 39: 1666-1672. <http://www3.interscience.wiley.com/cgi-bin/fulltext/120748701/pdfstart>.
- Emri, M., T. Marian, L. Tron, L. Balkay and Z. Krasznai, 1998. Temperature adaptation changes ion concentrations in spermatozoa and seminal plasma of common carp without affecting sperm motility. *Aquaculture*, 167: 85-94. DOI: 10.1016/S0044-8486(98)00309-3.

- Lahnsteiner, F., R.A. Patzner and T. Weismann, 1993. Energy resources of spermatozoa of the rainbow trout, *Oncorhynchus mykiss*. (Pisces, Teleostei). *Reprod. Nutr. Dev.*, 33: 349-360. http://md.edpsciences.org/index.php?Option=article&access=standard&Itemid=129&url=/articles/md/pdf/1993/04/RND_0926-5287_1993_33_4_ART0004.pdf.
- Lahnsteiner, F., R. Patzner and T. Weismann, 1994. The testicular main duct and the spermatic duct in some cyprinid fishes. II. Composition of the seminal fluid. *J. Fish Biol.*, 44: 459-467. DOI: 10.1111/j.1095-8649.1994.tb01226.
- Lahnsteiner, F., B. Berger, T. Weismann and R.A. Patzner, 1995. Fine structure and motility of spermatozoa and composition of the seminal plasma in the perch. *J. Fish Biol.*, 47: 492-508. DOI: 10.1111/j.1095-8649.1995.tb01917.
- Lahnsteiner, F., N. Mansour and B. Berger, 2004. Seminal plasma proteins prolong the viability of rainbow trout (*Oncorhynchus mykiss*) spermatozoa. *Theriogenology*, 62: 801-808. DOI: 10.1016/j.theriogenology.2003.12.001.
- Linhart, O., S.D. Mims, B. Gomelsky, A.E. Hiott, W.L. Shelton, J. Cosson, M. Rodina and D. Gela, 2000. Spermiation of paddlefish (*Polyodon spathula*) stimulated with injection of LHRH analogue and carp pituitary extract. *Aquat. Living Resour.*, 13: 455-460. DOI: 10.1016/S0990-7440(00)01068-8.
- Linhart, O., M. Rodina, D. Gela and M. Kocour, 2004. Optimization of artificial propagation in European catfish, *Silurus glanis* L. *Aquaculture*, 235: 619-632. DOI: 10.1016/j.aquaculture.2003.11.031.
- Morisawa, M., K. Suzuki, H. Shimizu, S. Morisawa and K. Yasuda, 1983. Effects of osmolality and potassium on motility of spermatozoa from freshwater cyprinid fishes. *J. Exp. Biol.*, 107: 95-103. <http://jeb.biologists.org/cgi/reprint/107/1/95?Maxtoshow=&HITS=10&hits=10&RESULTFORMAT=&andorexacttitle=&andorexacttitleabs=&andorexactfulltext=&searchid=1&FIRSTINDEX=0&sortspec=relevance&volume=107&firstpage=95&fdate=10/1/1980&tdate=9/30/1984&resourcetype=hwcit>.
- Piironen, J. and H. Hyvarinen, 1983. Composition of milt of some teleost fishes. *J. Fish Biol.*, 22: 351-361. DOI: 10.1111/j.1095-8649.1983.tb04757.
- Secer, S., N. Tekin, Y. Bozkurt, N. Bukan and E. Akcay, 2004. Correlation between biochemical and spermatological parameters in rainbow trout (*Oncorhynchus mykiss*) semen. *Isr. J. Aquac. Bamidgeh*, 56 (4): 274-280. <http://www.siamb.org.il/uploads/564Bozkurtabs.pdf>.
- Soengas, J.L., B. Sanmartin, P. Barciela, M. Aldegunde, G. Rozas, 1993. Changes in carbohydrate metabolism in domesticated rainbow trout *Oncorhynchus mykiss* related to spermatogenesis. *Comp. Biochem. Physiol.*, 105: 293-301. DOI: 10.1016/0300-9629(93)90211-L.
- Tekin, N., S. Secer, E. Akcay, Y. Bozkurt and S. Kayam, 2003. The effect of age on spermatological properties in rainbow trout (*Oncorhynchus mykiss* W. 1792). *Turk. J. Vet. Anim. Sci.*, 27 (1): 37-44. <http://journals.tubitak.gov.tr/veterinary/issues/vet-03-27-1/vet-27-1-6-010627.pdf>.
- White, I. and J. Macleod, 1963. Composition and Physiology of Semen. In: Hartman C.G. (Ed.). *Mechanisms Concerned with Conception*. Pergamon Press, London. LC, pp: 135-172. QP273 C625 1959.