

SP006

**VASOPRESSIN-RELATED COPEPTIN IS A NOVEL PREDICTOR OF EARLY ENDOTHELIAL DYSFUNCTION IN PATIENTS WITH ADULT POLYCYSTIC KIDNEY DISEASE**

Ismael Kocuyigit<sup>2</sup>, Mahmut Yilmaz<sup>3</sup>, Ozkan Gungor<sup>4</sup>, Eray Eroglu<sup>2</sup>, Aydin Unal<sup>2</sup>, Ozcan Orscelik<sup>6</sup>, Bulent Tokgoz<sup>2</sup>, Murat Sipahioglu<sup>2</sup>, Ahmet Sen<sup>1</sup>, Juan Carrero<sup>5</sup>, Oktay Oymak<sup>2</sup>, Jonas Axelsson<sup>5</sup>

<sup>1</sup>Biochemistry Erciyes University Faculty of Medicine Kayseri Turkey, <sup>2</sup>Nephrology Erciyes University Faculty of Medicine Kayseri Turkey, <sup>3</sup>Nephrology Gulhane School of Medicine Ankara Turkey, <sup>4</sup>Nephrology Kahramanmaraş Sutcu Imam University Faculty of Medicine Kahramanmaraş Turkey, <sup>5</sup>Division of Renal Medicine, Department of Clinical Science Karolinska Institutet Stockholm Sweden and <sup>6</sup>Cardiology Mersin University Faculty of Medicine Kayseri Turkey

**INTRODUCTION AND AIMS:** In this study, we examined the relative usefulness of serum copeptin levels as a surrogate marker of vasopressin (AVP) by correlating it with baseline and longitudinal changes in markers of both renal function and common CVD manifestations (hypertensive vascular disease, atherosclerosis and endothelial dysfunction) that accompany the progression of adult polycystic kidney disease (ADPKD).

**METHODS:** We studied a cohort of young and otherwise healthy ADPKD patients (n=235) and measured cardiovascular function using flow-mediation dilatation (FMD), carotid intima media thickness (cIMT) and pulse wave velocity (PWV), as well as serum copeptin (commercial ELISA, a stable marker of AVP activity). The same analyses were carried out at baseline and after three years of follow-up.

**RESULTS:** At baseline, median eGFR was 69 mL/min/1.73m<sup>2</sup>, mean FMD 6.9±0.9%, cIMT 0.7±0.1 mm, and PWV 8.1±1.2 m/s. At follow-up, equivalent values were 65 (44-75) mL/min/1.73m<sup>2</sup>, 5.8±0.9%, 0.8±0.1 mm, and 8.2±1.3 m/s. with all changes statistically significant. Plasma copeptin also rose from 0.62±0.12 to 0.94±0.19 ng/mL and this change correlated with ΔeGFR (-0.33, p<0.001), ΔFMD (0.599, p<0.001), ΔcIMT (0.562, p<0.001) and ΔPWV (0.27, p<0.001) also after linear regression modeling to correct for confounders (ΔAverage 24-h DBP, mmHg, ΔProteinuria, ΔcIMT, and ΔCRP). Finally, ROC analysis was done for a high copeptin with ΔeGFR [cut-off:<59], ΔFMD [cut-off:<7.08], ΔcIMT [cut-off:>0.76], ΔPWV, [cut-off:<7.80] and at baseline predicted future change in FMD and cIMT.

**CONCLUSIONS:** Vascular dysfunction as reflected by FMD and cIMT, but not PWV or an altered cardiac geometry, precede most other signs of disease in ADPKD but is predicted by elevated levels of the circulating AVP-marker copeptin.

ISSN 0931-0509 (Print) ISSN 1460-2385 (Online)  
Volume 32 Number S3 May 2017

# ndt

NEPHROLOGY DIALYSIS TRANSPLANTATION

Basic and clinical renal science

**54<sup>TH</sup> ERA-EDTA  
CONGRESS**  
Madrid, Spain  
June 3<sup>rd</sup>-6<sup>th</sup> 2017  
**ABSTRACTS**



Leading European Nephrology

June 3<sup>rd</sup>-6<sup>th</sup> 2017



[www.era-edta2017.org](http://www.era-edta2017.org)



Cristina Ortiz, "Moving forward", 2014,  
mixed technique, 70x120 cm, from the exhibition "Aspirations"

OXFORD  OPEN

OXFORD  
UNIVERSITY PRESS

[www.ndt.oxfordjournals.org](http://www.ndt.oxfordjournals.org)



Official Publication of the European  
Renal Association - European Dialysis  
and Transplant Association