



Head-to-head bisbenzazole derivatives as antiproliferative agents: design, synthesis, in vitro activity, and SAR analysis

Ronak Haj Ersa¹ · Mehmet Abdullah Alagoz² · Tugba Ertan-Boelli³ · Nizami Duran⁴ · Serdar Burmaoglu⁵ · Oztekin Algul¹


Received: 15 April 2020 / Accepted: 9 June 2020 / Published online: 18 June 2020
© Springer Nature Switzerland AG 2020

Abstract

In the present work, a series of bisbenzazole derivatives were designed and synthesized as antiproliferative agents. The antiproliferative activity of these compounds was investigated using MTT assay. Bisbenzazole derivatives showed significant antiproliferative activity against all the four tested cancer cell lines. Among the various bisbenzazole derivatives, bisbenzoxazole derivatives exhibited the most promising anticancer activity followed by bisbenzimidazole and bisbenzothiazole derivatives. All the derivatives were found to be less toxic as compared to methotrexate (positive control) in normal human cells, indicating selective and efficient antiproliferative activity of these bisbenzazole derivatives. The structure–activity relationships of heteroaromatic systems and linkers present in bisbenzazole derivatives were analyzed in detail. In silico ADMET prediction revealed that bisbenzazole is a drug-like small molecule with a favorable safety profile. Compound **31** is a potential antiproliferative hit compound that exhibits unique cytotoxic activity distinct from methotrexate.

Graphic abstract

Twenty-one bisbenzoxazole derivatives have been designed synthesized and evaluated to be an antiproliferative activity against four human tumor cell lines.



Keywords Bisbenzimidazole · Bisbenzoxazole · Bisbenzothiazole · Antiproliferative activity · SAR · ADMET

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11030-020-10115-0>) contains supplementary material, which is available to authorized users.

✉ Serdar Burmaoglu
serdarburmaoglu@atauni.edu.tr; sboglu@atauni.edu.tr

✉ Oztekin Algul
oztekinalgul@mersin.edu.tr

Extended author information available on the last page of the article

Introduction

Benzazoles are a family of heterocyclic compounds having a chemical skeleton consisting of a benzene ring fused with azole rings. Benz-fused azoles are among the most imperative class of molecules having a common heterocyclic

scaffold found in several biologically active and medicinally significant compounds.

These compounds are known to exhibit several therapeutic activities, including anticancer, antimicrobial, antiparasitic, antiviral, antihistamine, fungicidal, and antitubercular activity [1–10]. Some of the well-known drugs with benzazole rings that are used in clinical applications are omeprazole, emedastine, candesartan, astemizole, bezitramide, domperidone, lansoprazole, flunoxaprofen, and riluzole.

Bisbenzazoles consist of two benzazole nuclei fused together using a variety of linkers. The wide spectrum of pharmacological activities displayed by bisbenzazole derivatives makes them a highly important scaffold from drug development perspective. There are several reports on the medicinal activities/properties of this class of compounds. Various studies have revealed that bis-derivatives show better antiproliferative activity than the monomeric compounds [11]. Ueki et al. and Sato et al. demonstrated the cytotoxic activity of UK-1 and AJI95618 against B-16, HeLa, and P-338 cancer cell lines [12, 13]. Hoechst 33258 has undergone phase I clinical evaluation as an anticancer agent. It has been proposed that Hoechst 33258 acts via inhibition of topoisomerase and DNA helicase [14, 15]. Rance et al. demonstrated the antiproliferative activity of novel bisbenzothiazole analogs [16]. Among them, bisbenzazole moiety is one of the most important structures in drug discovery. A number of studies have been conducted in past that involved design and synthesis of novel bioactive bisbenzazole derivatives and evaluation of their antiproliferative activity against different human cancer cell lines (Fig. 1) [17–19].

All these studies indicated the therapeutic significance of bisbenzazole derivatives. In the present study, head-to-head bisbenzazole derivatives were designed and synthesized by accumulation of two benzazole units with different aliphatic and heteroaliphatic linkers with the aim to obtain/achieve better antiproliferative agents.

As shown in Fig. 2, the designing of bisbenzazole derivatives with different linkers was broadly divided into two segments. The first segment involves main backbone






Fig. 2 Molecular design strategy for the synthesis novel bisbenzazole derivatives connected via aliphatic and heteroaliphatic linkers


Fig. 1 Bisbenzazole scaffolds as antiproliferative agents



of the design, a benzazole unit that acts as a source of hydrogen-bond acceptor and donor. This would help to enhance the pharmacophoric properties as they exhibit drug-like properties. The second segment included aliphatic linkages with or without heteroatoms, which were generally used to control the lipophilicity and flexibility of the skeleton, and in silico ADMET prediction was also performed using the compounds.



Scheme 1 Synthesis 2-((1H-benzazol-2-yl)thio)acetic acid (5–7) from 2-mercaptobenzazoles (1–3) and chloroacetic acid (4)



Scheme 2 Synthesis of symmetric and asymmetric bisbenzazole derivatives using PPA method (20–37)

Results and discussion

Chemistry

The symmetric and asymmetric bisbenzazole derivatives (17–37) were synthesized using one-step and multi-step synthetic pathways as illustrated in Schemes 1 and 2. Polyphosphoric acid (PPA) method was used for the synthesis of desired compounds (20–37) (Scheme 2). For the synthesis of asymmetric compounds (17–19), 2-((1H-benzazol-2-yl)thio)acetic acid derivatives (5–7) were used as the starting material. These were prepared from 2-mercaptobenzazoles (1–3) and chloroacetic acid, followed by treatment with potassium hydroxide (Scheme 2).

The structures of all the designed and synthesized bisbenzazole derivatives (17–37) are shown in Scheme 2. All synthetic analogs (17–37) were characterized by physical and spectral analysis (FTIR, ¹H NMR, ¹³C NMR) and

elemental analysis. (For details, please see the Supplementary file.)

The chemical shifts of the ^1H NMR signals of bisbenzazole derivatives were investigated, and the signal of two protons of $-\text{CH}_2-$ linker group and four protons of $-\text{CH}_2-\text{CH}_2-$ linker group between benzazole rings is observed as a singlet with chemical shift values 3.35–4.88 ppm, while the signal of six protons of $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ linker group between benzazole rings is observed as a triplet of four protons of two terminals $-\text{CH}_2-$ attached to benzazole rings with chemical shift values 2.88–3.21 ppm and pentet or multiplet of two protons of $-\text{CH}_2-$ center of symmetry with chemical shift values 2.26–2.45 ppm. The signal of two protons of $-\text{CH}_2-\text{S}-$ linker group and four protons of $-\text{CH}_2-\text{S}-\text{CH}_2-$, $-\text{CH}_2-\text{NH}-\text{CH}_2-$, $-\text{CH}_2-\text{O}-\text{CH}_2-$ linker groups between benzazole rings is observed as a singlet with chemical shift values 3.58–5.20 ppm. The signal of aromatic $-\text{NH}-$ protons of bisbenzimidazole rings is observed as a singlet with chemical shift values 12.70–13.69 ppm. The signal of aromatic protons of benzazole rings is observed as (s, d, dd, ddd, td, t, and m) with chemical shift values 6.87–8.17 ppm. The chemical shifts of the ^{13}C NMR signals of bisbenzazole derivatives were investigated; the values were 22.6–69.9 ppm for aliphatic carbons and 169.3–109.3 ppm for aromatic carbons.

Antiproliferative Activity

The in vitro antiproliferative activities of all the synthesized compounds (**17–37**) were evaluated in four human cancer cell lines, lung cancer (A549), kidney cancer (A498), cervical cancer (HeLa), and liver cancer (HepG2), using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT assay) [20–22]. An antiproliferative drug, methotrexate, was used as the standard for comparison. The results of the MTT assay are summarized in Table 1. The antiproliferative activity of each analog was presented as the concentration of the compound that led to a 50% inhibition (IC_{50}) of cancer cell growth. These compounds were further screened for their cytotoxic nature using Vero cells (normal cell line). This was used to establish the selectivity of these active compounds toward cancerous cells. A potent anticancer compound should be more selective toward cancer cells and less toxic toward the normal cells.

To assess the specificity of the compounds, their toxicity levels were tested against Vero cells. The specificity of the compounds was calculated as their IC_{50} values for normal cells divided by their IC_{50} values for the specific cancer cells. Table 1 shows the specificity of all compounds.


Additionally, antiproliferative activity studies were performed at non-toxic concentrations of the compounds that were determined on the Vero cell line. The most specific

compound **31** and MTX were exposed for a period of 0–96 h, and A549 cells were evaluated and subjected to an assessment of their cytotoxicity responses using an MTT assay (Fig. 3). The proliferation of cancer cells treated with a non-toxic and lower-concentration dose of the compound was inhibited in a time-dependent manner. After 24, 48, and 72 h of treatment, compound **31** was more effective, and the number of live cells was lower than was observed with MTX. The antiproliferative activity of the compound increased as the exposure time was extended, and after 96-h treatment, the tested compounds had an activity equal to that of MTX.

As shown in Table 1, all the synthesized compounds showed significant antiproliferative activity against all the tested cancer cell lines. Among the synthesized compounds, bisbenzimidazole compounds **20**, **21**, and **23**; bisbenzoxazole compounds **18**, **27**, **30**, and **31**; and bisbenzothiazole compounds **32** and **33** showed promising antiproliferative activity against all the tested cancer cell lines. Among the asymmetric compounds bisbenzoxazole **18** and among the symmetric compounds bisbenzoxazole derivatives **30** and **31**, and bisbenzothiazole **32** were found to be most potent as antiproliferative agents and showed low toxicity and high specificity.

As most of the compounds from this series showed potency against tested human cancer cell lines, they can serve as attractive lead molecules for the discovery of novel antiproliferative agents in future. When the compounds were evaluated according to selectivity, for A549 cells, **18**, **30**, and **31** were found to be most potent antiproliferative agents with IC_{50} values of 0.28, 0.28, and 0.56 μM , respectively. The compounds **18**, **30**, **32**, and **33** showed good antiproliferative activities against both A498 and HeLa cells. For these two cell lines, **18**, **30**, and **33** displayed IC_{50} values of 0.28 μM , 0.56 μM , and 0.52 μM , respectively. Among the various compounds screened, **27** showed better specificity against A498, HeLa, and HepG2 cell lines. For HepG2 cells, **30** and **33** were found to be most potent as indicated by IC_{50} values of 0.28 μM . In addition to this, the compounds **18**, **30**, and **33** displayed potency against all the four cell lines. The compounds **18**, **30**, and **33** showed IC_{50} values of 0.28, 0.28, and 0.68 μM , respectively, in A549; IC_{50} values of 0.28, 0.56, and 0.52 μM , respectively, in A498; IC_{50} values of 0.56, 0.56, and 0.56 μM , respectively, in HeLa; and IC_{50} values of 0.56, 0.28, and 0.28 μM , respectively, in HepG2 cells.

For cytotoxicity test in Vero cells, **31** showed IC_{50} value of 17.90 μM , indicating 32 times more selectivity toward A549 and eight times more selectivity toward A498 and HeLa cancer cell lines. Similarly, **27** displayed IC_{50} value of 18.92 μM against Vero cells. Thus, **27** was found to be 16 times more selective toward A498, HeLa, and HepG2 cells and eight times more selective toward A549 cancer cell line.

Table 1 In vitro antiproliferative activity of synthesized bisbenzazole compounds studied using MTT assay

Compound no.	A	X	IC ₅₀ (μM) ^a				Specificity (SI) ^d					
			Vero ^b		HeLa ^c		HepG2 ^c		HeLa		HepG2	
			A549 ^c	A498 ^c	HeLa ^c	HepG2 ^c	A549	A498	HeLa	HepG2		
17	-NH-	-S-CH ₂ -	27.66±2.19	221.38±10.7	221.38±5.36	442.70±5.12	221.38±7.06	0.12	0.12	0.06	0.12	
18	-O-	-S-CH ₂ -	1.13±0.84	0.28±0.12	0.28±0.63	0.56±0.79	0.56±0.41	4.00	4.00	2.00	2.00	
19	-S-	-S-CH ₂ -	12.44±0.86	48.60±2.24	49.52±2.57	49.35±2.81	49.24±3.56	0.26	0.25	0.25	0.25	
20	-NH-	-CH ₂ -	5.030±0.07	1.26±1.55	1.26±0.07	2.52±1.77	2.52±0.78	4.01	4.01	2.00	2.00	
21	-NH-	-CH ₂ -CH ₂ -	0.60±0.25	0.60±0.38	1.19±0.27	0.60±0.29	0.60±0.13	1.00	0.50	1.00	1.00	
22	-NH-	-CH ₂ -CH ₂ -CH ₂ -	36.19±1.17	18.10±2.22	36.19±2.88	36.18±2.65	36.19±2.43	2.00	1.00	1.00	1.00	
23	-NH-	-CH ₂ -S-CH ₂ -	2.12±0.22	2.12±0.45	1.06±0.15	2.12±0.54	2.12±0.01.89	1.00	2.00	1.00	1.00	
24	-NH-	-CH ₂ -O-CH ₂ -	17.970±3.41	8.98±1.35	4.49±1.02	8.98±1.56	4.49±0.96	2.00	4.00	2.00	4.00	
25	-NH-	-CH ₂ -NH-CH ₂ -	14.10±4.32	55.17±3.52	56.36±5.22	56.35±2.63	56.36±3.31	0.26	0.25	0.25	0.25	
26	-O-	-CH ₂ -	9.99±1.28	4.99±0.98	4.99±1.67	4.99±1.48	9.99±2.57	2.00	2.00	2.00	1.00	
27	-O-	-CH ₂ -CH ₂ -	18.92±3.67	2.37±0.85	1.18±0.66	1.18±0.29	1.18±0.45	8.00	16.03	16.03	16.03	
28	-O-	-CH ₂ -CH ₂ -CH ₂ -	35.93±2.98	35.93±1.88	17.97±2.33	17.96±2.53	35.93±2.49	1.00	2.00	2.00	1.00	
29	-O-	-CH ₂ -S-CH ₂ -	33.75±1.02	16.87±1.05	16.87±2.63	16.87±0.36	16.87±2.33	2.00	2.00	2.00	2.00	
30	-O-	-CH ₂ -O-CH ₂ -	2.23±0.62	0.28±0.37	0.56±0.25	0.56±0.36	0.28±0.10	8.01	4.01	4.01	8.01	
31	-O-	-CH ₂ -NH-CH ₂ -	17.90±0.71	0.56±0.38	2.24±0.84	2.24±0.45	2.24±0.63	31.90	8.00	8.00	8.00	
32	-S-	-CH ₂ -	4.42±0.83	1.11±0.52	0.56±0.96	1.11±0.32	1.11±0.06	4.01	8.01	4.01	4.01	
33	-S-	-CH ₂ -CH ₂ -	1.05±0.16	0.68±0.55	0.52±0.39	0.56±0.72	0.28±0.96	2.00	2.00	2.00	4.00	
34	-S-	-CH ₂ -CH ₂ -CH ₂ -	32.21±2.97	16.11±1.49	16.11±5.33	16.10±1.43	16.11±0.85	2.00	2.00	2.00	2.00	
35	-S-	-CH ₂ -S-CH ₂ -	7.61±1.04	7.61±1.56	3.81±0.44	3.81±0.36	7.61±1.79	1.00	2.00	2.00	1.00	
36	-S-	-CH ₂ -O-CH ₂ -	4.00±1.73	4.00±1.46	2.00±1.01	4.00±0.75	4.00±1.39	1.00	2.00	1.00	1.00	
37	-S-	-CH ₂ -NH-CH ₂ -	32.11±3.55	16.10±1.86	16.10±2.57	16.05±3.55	16.10±2.84	2.00	2.00	2.00	2.00	
Methotrexate ^e			0.030±0.41	0.025±0.53	0.016±0.28	0.022±0.36	0.041±0.12	1.20	1.87	1.36	0.73	

Abbreviation Vero; African green monkey kidney epithelial cell, A549; human lung adenocarcinoma epithelial cell line, A498; human renal cancer cell line HeLa; human cervical cancer cell line HepG2; human hepatocellular cell line. ^aThe reported values represent the mean±SD for each compound based on three independent experiments, ^bnormal kidney epithelial cell, ^ccancer cells, ^dthe SI values are calculated as the ratio of the IC₅₀ between the Vero's and cancer cells, ^eused as reference




Fig. 3 MTT assay on A549 cell line after 96 h with MTX and compound **31**. The absorbance values were selected as 570 nm for the MTT method. Control cells not containing compounds and MTX were incubated same conditions. Cell viability was calculated as the ratio of absorbance of treated cells with compound or MTX to untreated cells. Given values show the mean standard deviations from three independent experiments carried out in triplicate. *Note:* *p* values: 24 h: control—31; *p* < .000; control—MTX; *p* < .000; control—MTX *p* < .05. 48 h: control—31; *p* < .000; control—MTX; *p* < .000; control—MTX *p* < .05. 72 h: control—31; *p* < .000; control—MTX; *p* < .000; control—MTX *p* < .05. 96 h: control—31; *p* < .000; control—MTX; *p* < .000; control—MTX. NS: nonsignificant

Furthermore, compound **30** with IC_{50} value 2.23 μ M against Vero cells showed eight times more selectivity toward A549 and HepG2. The results of this comparative analysis suggested that the most potent compounds from this newly synthesized series were characterized by more selectivity toward cancerous cells and very less toxicity toward the normal cells.

SAR analysis

Most of the compounds from the series displayed moderate to good antiproliferative activity toward cancer cell lines. A careful examination of the data helped to establish a significantly regular SAR. The synthesized compounds were analogs of bisbenzoxazoles with varying alkyl or heteroalkyl linkers. Accordingly, SAR was established by the comparison of the antiproliferative activities of the different linker and bisbenzazole groups. Among these compounds, bisbenzimidazole derivatives **20**, **21**, and **23**; bisbenzoxazole derivatives **18**, **26**, **27**, **30**, and **31**; and bisbenzothiazole derivatives **32** and **33** exhibited the most potent antiproliferative activity. In contrast, compounds **22**, **24**, **25**, **28**, **29**, and **34–37** with different linker and bisbenzazole groups showed the lowest antiproliferative activity against all cancer cells. In particular, the 1C linker in bisbenzothiazole **32** and 2C linker in bisbenzimidazole and bisbenzothiazole **21** and **33** were found to be associated with increased potency of the compounds, as compared to the

bisbenzoxazoles **26** and **27**, respectively. In addition, the three-atom linker groups ($-CH_2-CH_2-CH_2-$, $-CH_2-O-CH_2-$, and $-CH_2-NH-CH_2-$) in the benzazole derivatives (**22–25**, **28**, **29**, and **34–37**) did not significantly improve the activities against cancer cell lines, except for compound **31** against A549 and compound **30** in all cancer cell lines. In general, the 3C linkers in all benzazole scaffolds were inactive compounds for antiproliferative activity.

The resulting data are presented in Table 1. It shows that all the active compounds had generally considerable activity against cancer cell lines with IC_{50} values of 0.28–2.12 μ M. Compound **30** was the most active agent against A549 and HepG2 cells with an IC_{50} value of 0.28 μ M.

Physicochemical and toxicology properties

After completing structural characterization and biological activity studies, important points to know about these compounds were their physicochemical and toxicological properties. (Details are provided in supplementary content of Table S1.) All compounds were analyzed using PRE-ADMET and DATAWARRIOR 4.07.02 software. (Details are provided in supplementary content of Table S2.) With respect to the toxicological parameters, all compounds were evaluated as good, as none showed the potential to be carcinogenic or mutagenic, and they showed a medium-level ability to inhibit the hERG potassium channel and CYP450. For ADME properties, in general, the compounds' inhibitors were well evaluated; only compound **31** presented a weak binding to plasma proteins, low absorption in the brain–blood barrier, and good absorption in the intestinal system. In the case of drug-like parameters, almost the all of the compounds reached values allowing them to be considered as potential oral drugs.

All of the synthesized compounds were found to comply with Lipinski's rule of five. With respect to the toxicological parameters, none of the synthesized compounds had estimated mutagenic, carcinogenic, irritant, or reproduction effects. In the case of drug-like parameters, all compounds can be considered as potential oral drugs because they are well absorbed in the human intestine. Except for compounds **29** and **35**, all compounds inhibit the hERG potassium channel. Only **25** of the compounds does not inhibit CYP450. In addition, compound **25**, unlike most of the others, follows the lead-like rule and is weakly bound to plasma proteins. (Details are provided in supplementary content of Table S2.)

Conclusion

Bisbenzazole analogs (**17–37**) were synthesized, characterized, and evaluated for their antiproliferative activities against different cell lines (A549, A498, HeLa, and HepG2). Having a $-\text{CH}_2-\text{O}-\text{CH}_2-$ linker at the 2-position of the benzoxazole moiety enhanced the cytotoxic activity against the designated cell lines at low micromolar concentrations. The structure–activity relationships of heteroaromatic systems and linkers present in bisbenzazole derivatives were analyzed in detail.

Compounds **18**, **21**, **30**, and **33** displayed the highest antiproliferative activity and lower IC_{50} (μM) values against all cancer cell lines, compared to MTX. More remarkably, compound **31** showed approximately 16-fold better cytotoxicity on A549 cells, with approximately 13-fold high specificity, compared to MTX. The results also demonstrated that compound **27** has significant antiproliferative effects on A498, HeLa, and HepG2 cells, in comparison with MTX. Moreover, compound **27** showed more selective but much higher IC_{50} values than other active compounds and MTX; therefore, it was a less active compound. As a result, we can conclude that the linker group consists of two or three atoms and that one of the atoms should be a heteroatom.

The clinical application of most anticancer drugs has been found to be associated with certain side effects or toxicity. These limitations of anticancer drugs are mostly attributed to the poor selectivity of drugs toward cancer cells. The selectivity of drugs is an important factor affecting their utility. In the present study, a pharmacophore hypothesis was developed to analyze SARs between the molecular structures of the synthesized bisbenzazole derivatives and observed biological activity in the A549 cell line. At the same time, the physicochemical properties of the compounds play an important role in reaching the active region of the cell and the interactions with the active region.

In conclusion, our studies have shown that bisbenzazole scaffolds have good antiproliferative effects on all cancer cell lines. However, linkers are also important for changing antiproliferative activities. SAR studies of symmetrical derivatives showed that 1C linker compounds have moderate activity, although 2C linker compounds displayed more improved antiproliferative activity against all tested cancer cell lines. Interestingly, all three-atom linker compounds were inactive. Also, in general, three-atom linker compounds are less active than 2C linker compounds, except for compound **30**, which showed good antiproliferative activity against all cancer cell lines.

For asymmetrical bisbenzoxazole derivative, compound **18** exhibited good antiproliferative activity compared to other benzazoles (**17** and **19**).

Finally, in silico ADMET prediction highlighted that all compounds have desirable drug-like properties, favorable safety profiles, and comply with Lipinski's rule of five.

Experimental

Chemistry

All reagents used were commercially available unless otherwise specified, and all solvents were distilled before use. The precursor (**5–7**) was synthesized using the reported method in the literature [23]. Melting points (mp) were determined with Mettler Toledo MP90 melting point device. General reaction visualization was achieved by thin-layer chromatography (TLC) purchased from Merck KGaA (silica gel 60 F254) based on Merck DC plates (aluminum based) by using UV light (254 nm). Chromatographic separations were carried out using silica gel 60 (Merck, 63–200 μm particle size, 60–230 mesh). FTIR spectra were recorded as ATR on a Perkin Elmer Spectrum One FTIR spectrometer. All the nuclear magnetic resonance spectra of the analogs obtained on Bruker spectrometers (400 MHz for ^1H NMR and 100 MHz for ^{13}C NMR) in ppm (δ) refer to the solvent signal center at δ (7.19 and 76.0) ppm, δ (2.52 and 39.5) ppm, and δ (3.34 and 49.0) ppm for CDCl_3 , d_6 -DMSO, and d_4 - CH_3OH , respectively. Chemical shifts (δ) are reported in ppm. Coupling constants (J) are reported in Hz. Standard abbreviations indicating multiplicity were used as follows: br s (broad singlet), s (singlet), d (doublet), t (triplet), q (quartet), p (pentet), m (multiplet), and dd (doublet of doublets).

General procedure for preparation of compounds (20–37) 1,2-Phenylenediamine (**8**), 2-aminophenol (**9**), or 2-mercaptoanilin (**10**) (2 eq) and the corresponding dicarboxylic acid derivatives (**11**, **12**, **13**, **14**, **15**, and **16**) (1 eq) are heated for a period of 13–15 h in PPA at 180 °C. The reaction was monitored by thin-layer chromatography (TLC). UV (ultraviolet) light was used in the determination of stains in the works of TLC (Kieselgel 60 F254, ready-to-use aluminum plate coated with 0.2 mm thickness) which was made by using ready-made plates. After cooling, the reaction mixture was poured into ice water and neutralized by mixing with 5 M NaOH till slightly basic pH (8–9) to get the precipitate. The resulting precipitate was filtered off, washed with cold water, and crystallized with a suitable solvent. The resulting crystalline compounds were filtered, and the vacuumed product was dried.

Bis(1H-benzo[d]imidazol-2-yl)methane (20) The above procedure was followed with **8** and **11** to yield **20** as a brown powder solid (36% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.60; $mp = 119–123$ °C; IR (KBr, cm^{-1}) V_{max} 3058, 2962, 1656, 1314, 756, 672. The 1H NMR and ^{13}C NMR spectra are in agreement with the reported data [24, 25]. **Anal. calcd.** for $C_{15}H_{12}N_4$: C, 72.56; H, 4.87; N, 22.57. Found: C, 72.44; H, 4.95; N, 22.63.

1,2-Bis(1H-benzo[d]imidazol-2-yl)ethane (21) [26, 27] The above procedure was followed with **8** and **12** to yield **21** as a white powder solid (55% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.58; $mp = 119–123$ °C; IR (KBr, cm^{-1}) V_{max} 3387, 3200, 2719, 2608, 1626, 1574, 814, 764; 1H NMR (400 MHz, d_6 -DMSO) δ 7.84–7.76 (m, 4H, Ar–H), 7.60–7.49 (m, 4H, Ar–H), 3.91 (s, 4H, CH_2); ^{13}C NMR (100 MHz, d_6 -DMSO) δ 151.5, 131.3, 125.3, 113.8, 23.7. **Anal. calcd.** for $C_{16}H_{14}N_4$: C, 73.26; H, 5.38; N, 21.36. Found: C, 73.34; H, 5.29; N, 21.25.

1,3-Bis(1H-benzo[d]imidazol-2-yl)propane (22) The above procedure was followed with **8** and **13** to yield **22** as a brown powder solid (65% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.48; $mp = 270–275$ °C; IR (KBr, cm^{-1}) V_{max} 3048, 2952, 1547, 1435, 735. The 1H NMR and ^{13}C NMR spectra are in agreement with the reported data [28, 29]. **Anal. calcd.** for $C_{17}H_{16}N_4$: C, 73.89; H, 5.84; N, 20.27. Found: C, 73.74; H, 5.79; N, 20.25.

Bis((1H-benzo[d]imidazol-2-yl)methyl)sulfane (23) The above procedure was followed with **8** and **14** to yield **23** as a brown powder solid (47% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.71; $mp = 119–122$ °C; IR (KBr, cm^{-1}) V_{max} 3055, 2781, 1510, 1437, 725. The 1H NMR and ^{13}C NMR spectra are in agreement with the reported data [30–32]. **Anal. calcd.** for $C_{16}H_{14}N_4S$: C, 65.28; H, 4.79; N, 19.03; S, 10.89. Found: C, 65.14; H, 4.89; N, 19.15; S, 10.95.

2,2'-(Oxybis(methylene))bis(1H-benzo[d]imidazole) (24) The above procedure was followed with **8** and **15** to yield **24** as an orange powder solid (70% yield). The crystallization solvent is ethanol–water. R_f (chloroform/methanol 9:1) = 0.72; $mp = 122–126$ °C; IR (KBr, cm^{-1}) V_{max} 3056, 2915, 1439, 731. The 1H NMR spectrum is in agreement with the reported data [33, 34]. ^{13}C NMR (100 MHz, DMSO- d_6) δ 157.5, 151.3, 136.6, 134.3, 129.6, 127.8, 123.8, 121.6, 116.1, 114.8, 114.6, 62.5. **Anal. calcd.** for $C_{16}H_{14}N_4O$: C, 69.05; H, 5.07; N, 20.13. Found: C, 69.14; H, 4.97; N, 20.25.

Bis((1H-benzo[d]imidazol-2-yl)methyl)amine (25) The above procedure was followed with **1** and **16** to yield **25** as

a brown powder solid (47% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.60; $mp = 170–175$ °C; IR (KBr, cm^{-1}) V_{max} 3469, 3139, 2991, 1506, 1422, 747. The 1H NMR and ^{13}C NMR spectra are in agreement with the reported data [35–37]. **Anal. calcd.** for $C_{16}H_{15}N_5$: C, 69.29; H, 5.45; N, 25.25. Found: C, 69.32; H, 5.51; N, 25.37.

Bis(benzo[d]oxazol-2-yl)methane (26) The above procedure was followed with **9** and **11** to yield **26** as a brown powder solid (62% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.86; $mp = 121–123$ °C; IR (KBr, cm^{-1}) V_{max} 3093, 3066, 2956, 1614, 1570, 1239, 833, 742. The 1H NMR and ^{13}C NMR spectra are in agreement with the reported data [38, 39]. **Anal. calcd.** for $C_{15}H_{10}N_2O_2$: C, 71.99; H, 4.03; N, 11.19. Found: C, 72.12; H, 3.92; N, 11.15.

1,2-Bis(benzo[d]oxazol-2-yl)ethane (27) [40, 41] The above procedure was followed with **9** and **12** to yield **27** as an orange powder solid (60% yield). The crystallization solvent is ethanol–water. R_f (chloroform/methanol 95:05) = 0.59; $mp = 135–138$ °C; IR (KBr, cm^{-1}) V_{max} 3098, 3059, 2926, 1611, 1569, 1242, 831, 752. 1H NMR (400 MHz, $CDCl_3$) δ 7.66–7.57 (m, 2H, Ar–H), 7.47–7.38 (m, 2H, Ar–H), 7.26–7.23 (m, 4H, Ar–H), 3.51 (s, 4H, CH_2); ^{13}C NMR (100 MHz, $CDCl_3$) δ 163.9, 160.0, 140.2, 123.8, 123.2, 118.8, 109.4, 24.5. **Anal. calcd.** for $C_{16}H_{12}N_2O_2$: C, 72.72; H, 4.58; N, 10.60. Found: C, 72.61; H, 4.63; N, 10.49.

1,3-Bis(benzo[d]oxazol-2-yl)propane (28) [42, 43] The above procedure was followed with **9** and **13** to yield **28** as a pink powder solid (70% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.63; $mp = 146–150$ °C; IR (KBr, cm^{-1}) V_{max} 3051, 2997, 1571, 1450, 1240, 756; 1H NMR (400 MHz, $CDCl_3$) δ 7.62–7.54 (m, 2H, Ar–H), 7.43–7.36 (m, 2H, Ar–H), 7.26–7.20 (m, 4H, Ar–H), 3.06 (t, $J = 7.34$ Hz, 4H, $-CH_2$), 2.45 (p, $J = 7.34$ Hz, 2H, $-CH_2$); ^{13}C NMR (100 MHz, $CDCl_3$) δ 165.0, 149.9, 140.3, 123.6, 123.2, 109.3, 26.8, 22.6. **Anal. calcd.** for $C_{17}H_{14}N_2O_2$: C, 73.37; H, 5.07; N, 10.07. Found: C, 73.41; H, 4.98; N, 10.21.

Bis(benzo[d]oxazol-2-ylmethyl)sulfane (29) [44, 45] The above procedure was followed with **9** and **14** to yield **29** as an orange powder solid (55% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.50; $mp = 189–193$ °C; IR (KBr, cm^{-1}) V_{max} 3131, 2900, 1548, 1455, 1278, 744; 1H NMR (400 MHz, d_4 - CH_3OH) δ 7.78 (dd, $J = 1.44, 8.01$ Hz, 2H, Ar–H), 7.00 (td, $J = 1.53, 8.06$ Hz, 2H, Ar–H), 6.87 (dd, $J = 1.24, 8.08$ Hz, 2H, Ar–H), 6.87 (td, $J = 1.44, 8.03$ Hz, 2H, Ar–H), 3.63 (s, 4H, $-CH_2$); ^{13}C NMR (100 MHz, DMSO- d_6) δ 157.6, 130.9, 128.8,

128.6, 116.6, 35.4. **Anal. calcd.** for $C_{16}H_{12}N_2O_2S$: C, 64.85; H, 4.08; N, 9.45; S, 10.82. Found: C, 64.96; H, 4.03; N, 9.38; S, 10.95.

2,2'-(Oxybis(methylene))bis(benzo[d]oxazole) (30) [46] The above procedure was followed with **9** and **15** to yield **30** as a brown powder solid (35% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.50; **mp** = 191–194 °C; **IR** (KBr, cm^{-1}) V_{max} 3048, 2948, 1589, 1452, 1251, 795; 1H NMR (400 MHz, $CDCl_3$) δ 8.00 (d, J = 8.14 Hz, 2H, Ar–H), 7.83 (d, J = 7.94 Hz, 2H, Ar–H), 7.47 (t, J = 7.74 Hz, 2H, Ar–H), 7.36 (t, J = 7.19 Hz, 2H, Ar–H), 3.74 (s, 4H, $-CH_2$); ^{13}C NMR (100 MHz, $DMSO-d_6$) δ 159.7, 151.3, 129.3, 126.3, 125.0, 122.2, 116.6, 63.9. **Anal. calcd.** for $C_{16}H_{12}N_2O_3$: C, 68.56; H, 4.32; N, 9.99. Found: C, 68.65; H, 4.43; N, 10.11.

Bis(benzo[d]oxazol-2-ylmethyl)amine (31) The above procedure was followed with **9** and **16** to yield **31** as a red powder solid (30% yield). The crystallization solvent is ethanol–water. R_f (chloroform/methanol 95:05) = 0.25; **mp** = 132–136 °C; **IR** (KBr, cm^{-1}) V_{max} 3453, 3058, 2930, 1569, 1459, 1243, 783; 1H NMR (400 MHz, $CDCl_3$) δ 7.68 (dd, J = 3.35, 5.66 Hz, 2H, NH), 7.52–7.46 (m, 2H, Ar–H), 7.32 (t, J = 6.43 Hz, 4H, Ar–H), 3.58 (s, 4H, $-CH_2$); ^{13}C NMR (100 MHz, $CDCl_3$) δ 149.3, 127.0, 126.7, 123.5, 120.6, 116.7, 37.2. **Anal. calcd.** for $C_{16}H_{13}N_3O_2$: C, 68.81; H, 4.69; N, 15.05. Found: C, 68.75; H, 4.59; N, 15.17.

Bis(benzo[d]thiazol-2-yl)methane (32) The above procedure was followed with **10** and **11** to yield **32** as a green powder solid (49% yield). The crystallization solvent is ethanol–water. R_f (chloroform) = 0.33; **mp** = 170–174 °C; **IR** (KBr, cm^{-1}) V_{max} 3053, 2951, 1592, 1590, 1503, 1062, 854, 729. The 1H NMR and ^{13}C NMR spectra are in agreement with the reported data [24, 25]. **Anal. calcd.** for $C_{15}H_{10}N_2S_2$: C, 63.80; H, 3.57; N, 9.92; S, 22.71. Found: C, 63.71; H, 3.66; N, 9.81; S, 22.64.

1,2-Bis(benzo[d]thiazol-2-yl)ethane (33) The above procedure was followed with **10** and **12** to yield **33** as a yellow powder solid (55% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.37; **mp** = 137–140 °C; **IR** (KBr, cm^{-1}) V_{max} 3055, 2993, 1591, 1510, 1088, 877, 724. The 1H NMR and ^{13}C NMR spectra are in agreement with the reported data [47, 48]. **Anal. calcd.** for $C_{16}H_{12}N_2S_2$: C, 64.83; H, 4.08; N, 9.45; S, 21.64. Found: C, 64.95; H, 3.96; N, 9.56; S, 21.76.

1,3-Bis(benzo[d]thiazol-2-yl)propane (34) [49, 50] The above procedure was followed with **10** and **13** to yield **34** as a green powder solid (61% yield). The crystallization solvent is ethanol–water. R_f (chloroform) = 0.61; **mp** = 199–202 °C;

IR (KBr, cm^{-1}) V_{max} 3050, 2981, 1515, 1436, 1050, 760; 1H NMR (400 MHz, $CDCl_3$) δ 7.91 (d, J = 8.13 Hz, 2H, Ar–H), 7.77 (d, J = 7.99 Hz, 2H, Ar–H), 7.39 (t, J = 8.10, 2H, Ar–H), 7.29–7.27 (m, 2H, Ar–H), 3.21 (t, J = 7.05, 4H, $-CH_2$), 2.43 (m, 4H, $-CH_2$); ^{13}C NMR (100 MHz, $CDCl_3$) δ 170.6, 153.3, 135.2, 126.0, 124.8, 122.7, 121.5, 33.4, 29.0. **Anal. calcd.** for $C_{17}H_{14}N_2S_2$: C, 65.77; H, 4.55; N, 9.02; S, 20.66. Found: C, 65.85; H, 4.66; N, 9.17; S, 20.47.

Bis(benzo[d]thiazol-2-ylmethyl)sulfane (35) [51, 52] The above procedure was followed with **10** and **14** to yield **35** as a brown powder solid (49% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.54; **mp** = 189–193 °C; **IR** (KBr, cm^{-1}) V_{max} 3059, 2966, 1513, 1433, 1095, 762; 1H NMR (400 MHz, $CDCl_3$) δ 7.88 (d, J = 8.12 Hz, 2H, Ar–H), 7.76 (d, J = 7.95 Hz, 2H, Ar–H), 7.40–7.27 (m, 4H, Ar–H), 4.17 (s, 4H, $-CH_2$); ^{13}C NMR (100 MHz, $DMSO-d_6$) δ 155.9, 135.5, 135.1, 129.0, 128.7, 127.2, 125.8, 115.8, 33.6. **Anal. calcd.** for $C_{16}H_{12}N_2S_3$: C, 58.50; H, 3.68; N, 8.53; S, 29.29. Found: C, 58.63; H, 3.74; N, 8.67; S, 29.12.

2,2'-(Oxybis(methylene))bis(benzo[d]thiazole) (36) The above procedure was followed with **10** and **15** to yield **36** as a cream powder solid (67% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.72; **mp** = 103–106 °C; **IR** (KBr, cm^{-1}) V_{max} 3062, 2884, 1528, 143, 1038, 757. The 1H NMR spectrum is in agreement with the reported data [53, 54]. ^{13}C NMR (100 MHz, $DMSO-d_6$) δ 169.0, 152.6, 134.5, 126.3, 125.3, 122.7, 122.4, 69.9. **Anal. calcd.** for $C_{16}H_{12}N_2OS_2$: C, 61.51; H, 3.87; N, 8.97; S, 20.53. Found: C, 61.42; H, 3.73; N, 8.84; S, 20.44.

Bis(benzo[d]thiazol-2-ylmethyl)amine (37) The above procedure was followed with **10** and **16** to yield **37** as a yellow powder solid (40% yield). The crystallization solvent is ethanol–water. R_f (chloroform/methanol 95:05) = 0.27; **mp** = 101–104 °C; **IR** (KBr, cm^{-1}) V_{max} 3055, 2995, 1510, 1437, 1012, 724. The 1H NMR and ^{13}C NMR spectra are in agreement with the reported data [55, 56]. **Anal. calcd.** for $C_{16}H_{13}N_3S_2$: C, 61.71; H, 4.21; N, 13.49; S, 20.59. Found: C, 61.82; H, 4.19; N, 13.37; S, 20.48.

Synthesis of 2-((benzazol-2-yl)thio)acetic acid (5–7) A mixture of 2-mercaptobenzimidazole (**1**), 2-mercaptobenzoxazole (**2**), 2-mercaptobenzothiazole (**3**) (1 eq), and potassium hydroxide (1.2 eq) in methanol (20 mL) was stirred for 1 h. Then, 2-chloroacetic acid (**4**) (1.1 eq) was added into the mixture and the reaction mixture was refluxed for 8 h. After monitoring the reaction with TLC, the solvent was removed under reduced pressure, washed with cold water, dried, and recrystallized in ethanol to furnish the desired compounds (**5–7**).

2-((1H-benzo[d]imidazol-2-yl)thio)acetic acid (5) [23] The above procedure was followed with **1** and **4** to yield **5** as a white powder (91% yield). The crystallization solvent is ethanol. R_f (ethyl acetate/hexanes 2:1) = 0.16; mp = 120–123 °C; $^1\text{H NMR}$ (400 MHz, d_6 -DMSO) δ 7.46–7.44 (m, 2H, Ar–H), 7.15–7.12 (m, 2H, Ar–H), 4.15 (s, 2H, $-\text{CH}_2$).

2-(Benzo[d]oxazol-2-ylthio)acetic acid (6) [23] The above procedure was followed with **2** and **4** to yield **6** as a pink crystal (75% yield). The crystallization solvent is ethanol. R_f (ethyl acetate/hexanes 2:1) = 0.2; mp = 110–113 °C; $^1\text{H NMR}$ (400 MHz, d_6 -DMSO) δ 7.66–7.63 (m, 2H, Ar–H), 7.36–7.32 (m, 2H, Ar–H), 4.22 (s, 2H, $-\text{CH}_2$).

2-(benzo[d]thiazol-2-ylthio)acetic acid (7) [23] The above procedure was followed with **3** and **4** to yield **7** as a white powder (83% yield). The crystallization solvent is ethanol. R_f (ethyl acetate/hexanes 2:1) = 0.16; mp = 118–121 °C; $^1\text{H NMR}$ (400 MHz, d_6 -DMSO) δ 8.03 (d, J = 7.71 Hz, 1H, Ar–H), 7.87 (d, J = 7.14 Hz, 1H, Ar–H), 7.51–7.41 (m, 1H, Ar–H), 7.41–7.36 (m, 1H, Ar–H), 4.12 (s, 2H, $-\text{CH}_2$).

General Procedure for preparation of compounds (17–19) A mixture of *o*-phenylenediamine (**8**), 2-aminophenol (**9**), 2-mercaptoaniline (**10**) (1 eq), and 2-((benzazol-2-yl)thio)acetic acid (**5–7**) (1 eq) in PPA (5–7 g) was heated for 12 h in an oil bath at 150 °C. The reaction mixture was poured into ice water and neutralized by mixing with 5 M NaOH till slightly basic pH (8–9) to get the precipitate. The resulting precipitate was filtered off, washed with cold water, and recrystallized with ethanol–water.

2-(((1H-benzo[d]imidazol-2-yl)methyl)thio)-1H-benzo[d]imidazole (17) [57, 58] The above procedure was followed with **8** and **5** to yield **17** as a white powder (65% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.67; mp = 258–262 °C; IR (KBr, cm^{-1}) ν_{max} 3053, 2931, 1504, 1407, 849, 741. The $^1\text{H NMR}$ and $^{13}\text{C NMR}$ spectra are in agreement with the reported data. **Anal. calcd.** for $\text{C}_{15}\text{H}_{12}\text{N}_4\text{S}$: C, 64.26; H, 4.31; N, 19.98; S, 11.44. Found: C, 64.22; H, 4.19; N, 20.07; S, 11.58.

2-(Benzo[d]oxazol-2-ylmethyl)thio)benzo[d]oxazole (18) [57, 58] The above procedure was followed with **9** and **6** to yield **18** as a pink powder (56% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.63; mp = 146–150 °C; IR (KBr, cm^{-1}) ν_{max} 3093, 2995, 1501, 1445, 1238, 743; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.74–7.69 (m, 1H, Ar–H), 7.66–7.61 (m, 1H, Ar–H), 7.54–7.45 (m, 2H, Ar–H), 7.32 (m, 4H, Ar–H), 4.84 (s, 2H, $-\text{CH}_2$); $^{13}\text{C NMR}$ (100 MHz, $\text{DMSO}-d_6$) δ 156.1, 131.0, 115.9, 115.5, 115.3, 33.0. **Anal. calcd.** for $\text{C}_{15}\text{H}_{10}\text{N}_2\text{O}_2\text{S}$: C, 63.81; H,

3.57; N, 9.92; S, 11.36. Found: C, 63.92; H, 3.53; N, 10.01; O, S, 11.30.

2-(Benzo[d]thiazol-2-ylmethyl)thio)benzo[d]thiazole (19) [59] The above procedure was followed with **10** and **7** to yield **19** as a green powder (55% yield). The crystallization solvent is ethanol–water. R_f (ethyl acetate/hexanes 1:1) = 0.69; mp = 202–206 °C; IR (KBr, cm^{-1}) ν_{max} 3153, 2878, 1656, 1462, 1212, 768; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.01 (d, J = 8.16 Hz, 1H, Ar–H), 7.93 (d, J = 8.10 Hz, 1H, Ar–H), 7.82–7.75 (m, 2H, Ar–H), 7.50–7.29 (m, 4H, Ar–H), 5.05 (s, 2H, $-\text{CH}_2$); $^{13}\text{C NMR}$ (100 MHz, $\text{DMSO}-d_6$) δ 163.4, 161.0, 155.5, 138.4, 135.9, 130.5, 125.2, 116.0, 115.9, 115.8, 114.0, 113.8, 33.5. **Anal. calcd.** for $\text{C}_{15}\text{H}_{10}\text{N}_2\text{S}_3$: C, 57.29; H, 3.21; N, 8.91; S, 30.59. Found: C, 57.18; H, 3.28; N, 8.87; S, 30.44.

Biochemistry

Cell culture Studies

All the synthesized compounds (**17–37**) were evaluated for their in vitro antiproliferative activity against four human cancer cell lines by comparing the results with the standard antiproliferative drug, methotrexate (MTX). In vitro antiproliferative screening of titled compounds against human hepatocellular cell line (HepG2), human renal cancer cell line (A498), human lung adenocarcinoma epithelial cell line (A549), and human cervical cancer cell line (HeLa) were performed using MTT assay [22]. The obtained results of in vitro antiproliferative activities are summarized in Table 1. The selectivity of these compounds toward cancerous cells is evaluated against Vero (African green monkey kidney epithelial cell) to determine the non-toxic concentrations of the compounds on relatively healthy cells. All cell lines were obtained from the cell culture collections of Mustafa Kemal University. Cell incubations were done at 37 °C in 5% (v/v) CO_2 .

MTT cell viability assay

The cytotoxic activity of the compounds was determined by using MTT assay [20, 21]. The compounds were dissolved in dimethyl sulfoxide (DMSO) and then diluted to the appropriate concentrations in Dulbecco's modified Eagle's medium (DMEM). Concentration of DMSO was less than 1% in the culture medium. Subsequently, the cells were seeded at 104 cells/well in DMEM, supplemented with 10% fetal bovine serum (FBS), 100 units/mL penicillin G, and 100 $\mu\text{g}/\text{mL}$ streptomycin in each well of 96-well microculture plates. The cells were cultured at 37 °C for 24, 48, 72, and 96 h in an incubator containing 5% CO_2 .

After incubation, cells were treated with test compounds of appropriate concentrations for 24, 48, 72, and 96 h. After incubation period, 10 μ L MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide) (5 mg/mL) was added to each well and the plates were further incubated for 4 h. Then, the medium of each well was carefully removed and formazan crystals were dissolved in 100 μ L of DMSO. Absorbance was determined at 570 nm for each well using a microplate reader (Bioteck) [60].

The cell viability was expressed as percentage of the viable cells in each sample with respect to the control wells. Three independent experiments in triplicates were done for the determination of the growth inhibition of each compound. The IC_{50} values were calculated from concentration–response curves using the SPSS (SPSS Inc., Chicago) software.

Acknowledgements We thank the Scientific and Technological Research Council of Turkey (TUBITAK, Grant Number: 115S190) and Mersin University (Project Number: 2018-1-TP3-2911) for their financial support.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

- Spasov AA, Yozhitsa IN, Bugaeva LI (1999) Benzimidazole derivatives: spectrum of pharmacological activity and toxicological properties. *Pharm Chem J* 33:232–243. <https://doi.org/10.1007/BF02510042>
- Gupta M, Paul S, Gupta R (2009) General characteristics and applications of microwaves in organic synthesis. *Acta Chim Slov* 56:749–764
- Piscitelli F, Ballatore C, Smith A (2010) Solid phase synthesis of 2-aminobenzothiazoles. *Bioorg Med Chem Lett* 20:644–648. <https://doi.org/10.1016/j.bmcl.2009.11.055>
- Balakumar C, Kishore DP, Rao KV, Narayana BL, Rajwinder K, Rajkumar V, Rao AR (2012) Design, microwave-assisted synthesis and in silico docking studies of new 4H-pyrimido[2,1-b]benzothiazole-2-arylamino-3-cyano-4-ones as possible adenosine A2B receptor antagonists. *Indian J Chem* 51B:1105–1113
- Patil A, Ganguly S, Surana S (2008) A systematic review of benzimidazole derivatives as an antiulcer agent. *RJC* 1:447–460
- Narasimhan B, Sharma D, Kumar P (2012) Benzimidazole: a medicinally important heterocyclic moiety. *Med Chem Res* 21:269–283. <https://doi.org/10.1007/s00044-010-9533-9>
- Kohli P, Srivastava SD, Srivastava SK (2007) Synthesis and biological activity of mercaptobenzoxazole based thiazolidinones and their arylidenes. *J Chin Chem Soc* 54:1003–1010. <https://doi.org/10.1002/jccs.200700144>
- Sivakumar R, Pradeepchandran R, Jayaveera KN, Kumarnallasi-van P, Vijaijanand PR, Venkatnarayanan R (2011) Benzimidazole: an attractive pharmacophore in medicinal chemistry. *Int J Pharm Sci Res* 3:19–31
- Deb PK, Kaur R, Chandrasekaran B, Bala M, Gill D, Kaki VR, Akkinipalli RR, Mailavaram R (2014) Synthesis, anti-inflammatory evaluation, and docking studies of some new thiazole derivatives. *Med Chem Res* 23:2780–2792. <https://doi.org/10.1007/s00044-013-0861-4>
- Palmer FJ, Trigg RB, Warrington JV (1971) Benzothiazolines as antituberculous agents. *J Med Chem* 14:248–251. <https://doi.org/10.1021/jm00285a022>
- Hadden MK, Blagg BS (2008) Dimeric approaches to anti-cancer chemotherapeutics. *Anticancer Agents Med Chem* 8:807–816. <https://doi.org/10.2174/187152008785914743>
- Ueki M, Ueno K, Miyadoh S, Abe K, Shibata K, Taniguchi M, Oi S (1993) A novel cytotoxic metabolite from *Streptomyces* sp. 517-02. *J Antibiot* 46:1089–1094. <https://doi.org/10.7164/antibiotics.46.1089>
- Sato S, Kajiura T, Noguchi M, Takehana K, Kobayashi T, Tsuji T (2001) A new cytotoxic benzoxazole derivative produced by *Streptomyces* sp. *J Antibiot* 54:102–104. <https://doi.org/10.7164/antibiotics.54.102>
- Jenkins TC (2000) Targeting multi-stranded DNA structures. *Curr Med Chem* 7:99–115. <https://doi.org/10.2174/0929867003375551>
- Singh MP, Joseph T, Kumar S, Bathini Y, Lown JW (1992) Synthesis and sequence-specific DNA binding of a topoisomerase inhibitory analog of Hoechst 33258 designed for altered base and sequence recognition. *Chem Res Toxicol* 5:597–607. <https://doi.org/10.1021/tx00029a003>
- Racane L, Kraljevic PS, Ratkaj I, Stepanic V, Pavelic K, Tralic-Kulenovic V, Karminski-Zamola G (2012) Synthesis and antiproliferative evaluation of some new amidino-substituted bis-benzothiazolyl-pyridines and pyrazine. *Eur J Med Chem* 55:108–116. <https://doi.org/10.1016/j.ejmech.2012.07.005>
- Kumbhare RM, Dadmal T, Kosurkar U, Sridhar V, Rao JV (2012) Synthesis and cytotoxic evaluation of thiourea and *N*-bis-benzothiazole derivatives: a novel class of cytotoxic agents. *Bioorg Med Chem Lett* 22:453–455. <https://doi.org/10.1016/j.bmcl.2011.10.106>
- Gravatt GL, Baguley BC, Wilson WR, Denny WA (1994) DNA-directed alkylating agents. 6. Synthesis and antitumor activity of DNA minor groove-targeted aniline mustard analogs of pibenzimol (Hoechst 33258). *J Med Chem* 37:4338–4345. <https://doi.org/10.1021/jm00051a010>
- Shi Z, Zhao D, Huang Y, Du Y, Cao X, Gong Z, Zhao R, Li J (2012) Discovery, synthesis, and evaluation of small-molecule signal transducer and activator of transcription 3 inhibitors. *Chem Pharm Bull* 60:1574–1580. <https://doi.org/10.1248/cpb.c12-00745>
- McGahon AJ, Martin SJ, Bissonnette RP, Mahboubi A, Shi Y, Mogil RJ, Nishioka WK, Green DR (1995) The end of the (cell) line: methods for the study of apoptosis in vitro. *Methods in Cell Biology: Cell Death* 46:153–185. [https://doi.org/10.1016/s0091-679x\(08\)61929-9](https://doi.org/10.1016/s0091-679x(08)61929-9)
- Mosmann T (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 65:55–63. [https://doi.org/10.1016/0022-1759\(83\)90303-4](https://doi.org/10.1016/0022-1759(83)90303-4)
- van Meerloo J, Kaspers GJL, Cloos J (2011) Cell sensitivity assays: the MTT assay. *Methods Mol Biol* 731:237–245. https://doi.org/10.1007/978-1-61779-080-5_20
- Lan P, Romero FA, Wodka D, Kassick AJ, Dang Q, Gibson T, Cashion D, Zhou G, Chen Y, Zhang X, Zhang A, Li Y, Trujillo ME, Shao Q, Wu M, Xu S, He H, MacKenna D, Staunton J, Chapman KT, Weber A, Sebhat IK, Makara GM (2017) Hit-to-lead optimization and discovery of 5-((5-([1',1'-Biphenyl]-4-yl)-6-chloro-1 H-benzo [d] imidazol-2-yl) oxy)-2-methylbenzoic Acid (MK-3903): A novel class of benzimidazole-based activators of AMP-activated protein kinase. *J Med Chem* 60:9040–9052. <https://doi.org/10.1021/acs.jmedchem.7b01344>


24. Elagab HA, Alt HG (2015) Structure–property–relationship studies with ethylene polymerization catalysts of Ti, Zr and V containing heterocyclic ligands. *Inorgan Chim Acta* 437:26–35. <https://doi.org/10.1016/j.ica.2015.08.002>
25. Qian J, Zhang Y, Yin X (2012) The synthesis and study of bis(1-octylbenzimidazol-2-yl)alkane oil-soluble corrosion inhibitor. *Huaxue Tongbao* 75:88–91
26. Heinrich J, Koenig NF, Sobottka S, Sarkar B, Kulak N (2019) Flexible vs. rigid bis(2-benzimidazolyl) ligands in Cu(II) complexes: impact on redox chemistry and oxidative DNA cleavage activity. *J Inorg Biochem* 194:223–232. <https://doi.org/10.1016/j.jinorgbio.2019.01.016>
27. Yilmaz U, Kucukbay H (2016) Synthesis and characterization of novel phosphoramidates containing benzimidazole moiety. *Phosphorus, Sulfur Silicon Relat Elem* 191:140–143. <https://doi.org/10.1080/10426507.2015.1067209>
28. Inamdar SM, More VK, Mandal SK (2013) CuO nano-particles supported on silica, a new catalyst for facile synthesis of benzimidazoles, benzothiazoles and benzoxazoles. *Tetrahedron Lett* 54:579–583. <https://doi.org/10.1016/j.tetlet.2012.11.091>
29. Berends HP, Stephan DW (1984) Copper (I) and copper (II) complexes of biologically relevant tridentate ligands. *Inorgan Chim Acta* 93:173–178. [https://doi.org/10.1016/S0020-1693\(00\)88159-1](https://doi.org/10.1016/S0020-1693(00)88159-1)
30. Mao S, Shen K, Shi X, Wu H, Han X, Li C, Huang G (2018) Synthesis, crystal structure and biological activity of two binuclear Ag(I) complexes with bis-benzimidazole thioether ligands. *Inorgan Chim Acta* 471:82–90. <https://doi.org/10.1016/j.ica.2017.10.038>
31. Rao SS, Reddy CV, Dubey PK (2014) Synthesis of *N,N*-disubstituted bisbenzimidazolesulphides of Potential Pharmacological Interest. *JCHPS* 6:1199–1204
32. Xu Y, Wu H, Zhang H, Aderinto SO, Yang Z (2016) Synthesis, crystal structures, and DNA-binding studies of two silver (I) complexes with 1, 3-bis(1-ethylbenzimidazol-2-yl)-2-thiopropane. *J Coord Chem* 69:2988–2998. <https://doi.org/10.1080/00958972.2016.1218484>
33. Mao S, Shen K, Shi X, Xu Y, Wu H (2017) Two silver (I) complexes with bis (benzimidazole)-2-oxopropane ligands: syntheses, crystal structures and DNA binding studies. *APPI* 31:e3747. <https://doi.org/10.1002/aoc.3747>
34. Wu HL, Yun RR, Wang KT, Li K, Huang XC, Sun T (2010) Synthesis, crystal structure, and spectrum properties of Cobalt (II) complexes based on tridentate 1, 3-bis(benzimidazol-2-yl)-2-oxopropane ligand and derivative. *Z Anorg Allg Chem* 636:629–633. <https://doi.org/10.1002/zaac.200900298>
35. Feng R, Hou Y, Wu Z, Yang Y, Nie F (2015) Structures and fluorescent properties of Cadmium (II) complexes with 1D and 2D Structures based on tridentate benzimidazole ligands. *Z Anorg Allg Chem* 641:1918–1925. <https://doi.org/10.1002/zaac.201400511>
36. Kopel P, Wawrzak D, Langer V, Cihalova K, Chudobova D, Vesely R, Adam V, Kizek R (2015) Biological activity and molecular structures of bis(benzimidazole) and trithiocyanurate complexes. *Molecules* 20:10360–10376. <https://doi.org/10.3390/molecules200610360>
37. Chen B, Morlanes N, Adogla E, Takanabe K, Rodionov VO (2016) An efficient and stable hydrophobic molecular cobalt catalyst for water electro-oxidation at neutral pH. *ACS Catal* 6:4647–4652. <https://doi.org/10.1021/acscatal.6b01237>
38. Dauer D, Stalke D (2014) Heterocyclic substituted methanides as promising alternatives to the ubiquitous nacnac ligand. *Dalton Trans* 43:14432–14439. <https://doi.org/10.1039/C4DT01008F>
39. Kretsch J, Kreyenschmidt A, Herbst-Irmer R, Stalke D (2018) Alkali metal complexes based on bisheterocyclomethanide ligands. *Dalton Trans* 47:12606–12612. <https://doi.org/10.1039/C8DT01678J>
40. Kumar R, Selvam C, Kaur G, Chakraborti AK (2005) Microwave-assisted direct synthesis of 2-substituted benzoxazoles from carboxylic acids under catalyst and solvent-free conditions. *Synlett* 9:1401–1404. <https://doi.org/10.1055/s-2005-868509>
41. Shirakawa Y, Masuda T (2013) Coupling agents for rubber/carbon black and rubber compositions containing them. *PCT Int Appl WO* 2013015425
42. Cakir B, Ucucu U, Buyukbingol E, Abbasoglu U (1989) Benzoxazoles: bis-benzoxazole derivatives, synthesis, antifungal activities and QSARs. *J Faculty Pharm Gazi Uni* 6:15–21
43. Terao H, Ono Y, Ito Y, Isogai M, Hamada T, Imanishi, Tsunoda A (1991) Organic nonlinear optical device. *Japanese Kokai Tokkyo Koho JP* 03188425
44. Hao Y, Chen Y (2016) Excited-state intramolecular single and double proton transfer emission of 2,5-bis(benzoxazol-2-yl) thiophene-3,4-diol. *Dyes Pigme* 129:186–190. <https://doi.org/10.1016/j.dyepig.2016.03.002>
45. Hao Y, Zheng M, Chen Y (2014) A highly stable and water-soluble fluorescent dye for fluorescence imaging of living cells. *J Mater Chem B* 2:7369–7374. <https://doi.org/10.1039/C4TB01210K>
46. Zhou H, Wang L, Yin B (2004) Synthesis of bis(2-benzoxazolymethyl) ether by dry reaction under microwave irradiation. *Huaxue Shiji* 26:308–311
47. Alt H, Elagab H, Al-Humydi A (2011) Ethylene Polymerisation. *PCT Int Appl WO* 2011088990
48. Chakraborti AK, Selvam C, Kaur G, Bhagat S (2004) An efficient synthesis of benzothiazoles by direct condensation of carboxylic acids with 2-aminothiophenol under microwave irradiation. *Synlett* 5:851–855. <https://doi.org/10.1055/s-2004-820012>
49. Katritzky AR, Liang D, Fan W (1988) Bridged cyanine dyes. Part 2 [1].1-(*N*-methyl-2-benzothiazolylinio)-3-(*N*-methyl-2-benzothiazolylene) and 1-(*N*-methyl-4-pyridinio)-3-(*N*-methyl-4-pyridylene)cyclopenta-1,4-dienes with fused rings. *J Heterocycl Chem* 25:1315–1319. <https://doi.org/10.1002/jhet.5570250509>
50. Rai C, Braunwarth JB (1961) Synthesis of bisbenzothiazoles I. *J Org Chem* 26:3434–3436. <https://doi.org/10.1021/jo01067a100>
51. Strasser CE, Jongh LAD, Raubenheimer HG, Cronje S (2011) 2,2'-(Sulfanediyl)dimethylene)bis(1,3-benzothiazole). *Acta Cryst E* 67:o622. <https://doi.org/10.1107/S1600536811004478>
52. Sih JC, Graber DR (1983) 2-Mercapto-1,3-benzoxazole: a useful reagent for the preparation of symmetrical and unsymmetrical sulphides. *J Org Chem* 48:3842–3845. <https://doi.org/10.1021/jo00169a058>
53. Das SK, Mathur P (1999) Copper (II) complexes with bis thiazole based ligands: spectral, cyclic voltammetric and EPR studies. *Indian J Chem A* 38A:1277–1282
54. Ushenko IK (1952) α,γ -Epoxythiocarbonyl cyanines. *I Zh Obshch Khim* 22:711–715
55. Finn MG, Rodionov VO (2009) Ligands for copper-catalyzed azide-alkyne cycloaddition reactions. *PCT Int Appl WO* 2009038685
56. Buehrdel G, Beckert R, Herzigova P, Petrlikova E, Schuch D, Birkner E, Goerls H (2009) A new synthesis of push-pull pyrroles, their oxidation to stable 3H-pyrroles and an unexpected anellation reaction. *Eur J Org Chem* 20:3404–3412. <https://doi.org/10.1002/ejoc.200900295>
57. Akpa SJ, Say MV, Zoakouma SPR, Fante B, Sissouma D, Adjou A (2016) Synthesis of 2-(benzylthio) benzimidazole, 2-[(benzimidazol-2-yl)methylthio]benzimidazole and structural analogues against *Haemoncus contortus*. *AFR J Pharm Pharmacol* 10:670–680. <https://doi.org/10.5897/AJPP2016.4557>
58. Bouchouit M, Said M, Kara M, Bouacida S, Merazig H, Kacem-Chaouche N, Chibani A, Zouchoune B, Belfaitah A, Bouraiou

- A (2016) Synthesis, X-ray structure, theoretical investigation, corrosion inhibition and antimicrobial activity of benzimidazole thioether and their metal complexes. *Polyhedron* 119:248–259. <https://doi.org/10.1016/j.poly.2016.08.045>
59. Zubarovskii VM (1951) 2-(Hydroxymethyl)benzothiazole and its transformations. *Zh Obshch Khim* 21:2055–2064
60. Sabina XJ, Karthikeyan J, Velmurugan G, Tamizh MM, Shetty AN (2017) Design and in vitro biological evaluation of substituted

chalcones synthesized from nitrogen mustards as potent microtubule targeted anticancer agents. *New J Chem* 41:4096–4109. <https://doi.org/10.1039/C7NJ00265C>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Affiliations

Ronak Haj Ersan¹ · Mehmet Abdullah Alagoz² · Tugba Ertan-Bolelli³ · Nizami Duran⁴ · Serdar Burmaoglu⁵  · Oztekin Algul¹ 

¹ Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Mersin University, 33169 Mersin, Turkey

² Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Inonu University, 44280 Malatya, Turkey

³ Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Ankara University, 06560 Ankara, Turkey

⁴ Department of Medical Microbiology, Medical Faculty, Mustafa Kemal University, 31100 Antakya, Hatay, Turkey

⁵ Department of Chemistry, Faculty of Science, Ataturk University, 25240 Erzurum, Turkey