

## ORIGINAL ARTICLE

## Antioxidative effects of cysteamine, hyaluronan and fetuin on post-thaw semen quality, DNA integrity and oxidative stress parameters in the Brown Swiss bull

S. Sariözkan<sup>1,2</sup>, P. B. Tuncer<sup>3</sup>, S. Büyükleblebici<sup>4</sup>, M. N. Bucak<sup>5</sup>, F. Cantürk<sup>6</sup> & A. Eken<sup>7</sup>

1 Faculty of Veterinary Medicine, Department of Reproduction and Artificial Insemination, Erciyes University, Kayseri, Turkey;

2 Genome and Stem Cell Center-GENKOK, Erciyes University, Kayseri, Turkey;

3 Republic of Turkey Ministry of Food, Agriculture and Livestock, General Directorate of Food and Control, Ankara, Turkey;

4 Faculty of Veterinary Medicine, Department of Reproduction and Artificial Insemination, Aksaray University, Aksaray, Turkey;

5 Faculty of Veterinary Medicine, Department of Reproduction and Artificial Insemination, Selcuk University, Konya, Turkey;

6 Faculty of Medicine, Department of Basic Sciences, Erciyes University, Kayseri, Turkey;

7 Faculty of Pharmacy, Department of Pharmaceutical Toxicology, Erciyes University, Kayseri, Turkey

### Keywords

Antioxidant activities—cryopreservation—DNA damage—lipid peroxidation—sperm parameters

### Correspondence

Serpil Sariözkan, PhD, Assoc. Prof. of Reproduction, Faculty of Veterinary Medicine, Department of Reproduction and Artificial Insemination, Erciyes University, Kayseri, Turkey.

Tel.: +90 352 3399484;

Fax: +90 352 3372740;

E-mail: sariozkan75@yahoo.com

Accepted: December 12, 2013

doi: 10.1111/and.12236

### Summary

The aim of this study was to compare the effectiveness of antioxidants including cysteamine (2.5, 7.5 mM), hyaluronan (0.25, 1 mg ml<sup>-1</sup>) and fetuin (5, 10 mg ml<sup>-1</sup>) in the freezing of Brown Swiss bull semen. The best percentages of CASA motilities were achieved with 10 mg ml<sup>-1</sup> of fetuin and 2.5 mM of cysteamine. For sperm morphology, 10 mg ml<sup>-1</sup> of fetuin and 2.5 mM of cysteamine had better protective effects ( $P < 0.001$ ). The results of hypo-osmotic swelling test showed that the percentage values of membrane integrity in all the groups, excluding that supplemented with 5 mg ml<sup>-1</sup> of fetuin, were higher than those of the control group ( $P < 0.001$ ). Results obtained for the DNA damage of sperm cells demonstrated that 0.25 mg ml<sup>-1</sup> of hyaluronan, and 2.5 and 7.5 mM of cysteamine led to lower rates of spermatozoa with damaged DNA, compared with the control group ( $P < 0.001$ ). The maintenance of superoxide dismutase and glutathione peroxidase antioxidant activities following freeze-thawing with 2.5 and 7.5 mM of cysteamine and 10 mg ml<sup>-1</sup> of fetuin was demonstrated to be at a higher level in comparison with the control group ( $P < 0.001$ ). Malondialdehyde formation was found to be lower in the groups supplemented with 0.25 mg ml<sup>-1</sup> of hyaluronan and 7.5 mM of cysteamine after the freeze-thawing process ( $P < 0.001$ ).

### Introduction

Cryopreservation procedures lead to cold shock, ice crystal formation, oxidative stress, osmotic changes and lipid-protein reorganisations within the cell membrane, which affect normal sperm functions and result in loss of motility, viability, fertilising ability, deterioration of acrosomal and plasma membrane integrity and structural damage to DNA (Watson, 1995; Aitken *et al.*, 1998; Bailey *et al.*, 2000; Vishwanath & Shannon, 2000; Medeiros *et al.*, 2002).

Oxidative stress is known as one of the major causes of adverse effects on sperm physiology, through the induction

of peroxidation in the sperm plasma membrane. The detrimental effect of oxidative stress on spermatozoa is known to arise from the generation of reactive oxygen species (ROS) (Aitken & Fisher, 1994; Sharma & Agarwal, 1996; Agarwal *et al.*, 2008). ROS are physiologically generated by spermatozoa. The physiological levels of ROS play a crucial role in reproductive processes such as sperm–oocyte interactions (De Lamirande *et al.*, 1997), implantation and early embryonic development (Sakkas *et al.*, 1998), sperm capacitation, acrosome reaction, maintenance of fertilising ability, and stabilisation of the mitochondrial capsule in the mid-piece in cattle (Agarwal *et al.*, 2008; Desai *et al.*, 2009; Gonçalves *et al.*, 2010).

However, an imbalance between ROS generation and scavenging activity is known as oxidative stress (Agarwal *et al.*, 2003), which triggers the loss of membrane integrity, enzyme inactivation, DNA damage and cell death and finally links with infertility (Halliwell, 1994; Sharma & Agarwal, 1996).

Reactive oxygen species cause DNA base oxidation (Dizdaroglu *et al.*, 1991; Jaruga *et al.*, 1994) and DNA strand breaks (Hughes *et al.*, 1996; Twigg *et al.*, 1998a). The transmission of genetic information to future generations depends on sperm DNA integrity. DNA integrity has become an important indicator of sperm fertility (Hughes *et al.*, 1999). Excessive ROS generation has been associated with impaired DNA integrity as well as other impaired sperm quality parameters, which in return decrease the capability of the spermatozoa to fuse with the oocyte and thereby result in infertility (Aitken *et al.*, 1991; Sharma & Agarwal, 1996; Potts *et al.*, 2000). Spermatozoa contain various antioxidants associated with ROS scavenging, such as superoxide dismutase (SOD) and glutathione peroxidase (GPx) (Geva *et al.*, 1996; Garrido *et al.*, 2004). GPx is a selenium-containing antioxidative enzyme, which plays a major role in scavenging excessive free radicals against lipid peroxidation attack and acts directly as an antioxidant. Owing to its vital role in sperm maturation, the lack of GPx may reduce fertilising capacity (De Lamirande & Gagnon, 1993; Vernet *et al.*, 1997). Another ROS scavenger found in seminal plasma is SOD, which is derived primarily from the prostate gland and helps prevent oxidative damage to spermatozoa (Ball *et al.*, 2000; Baumber & Ball, 2005). Malondialdehyde (MDA) is an indicator of oxidative stress and end product of lipid peroxidation. High levels of MDA cause detrimental damage to sperm functions (Geva *et al.*, 1996). As demonstrated in recent studies, the addition of various antioxidants to the semen extender protects spermatozoa against the detrimental effects of ROS and improves post-thaw sperm functions such as motility, viability and fertility (Bilodeau *et al.*, 2001; Pena *et al.*, 2004; Sariözkan *et al.*, 2009; Bucak *et al.*, 2010a).

Cysteamine belongs to a large class of antioxidants known as thiol compounds. It increases the synthesis of both glutathione and other potent antioxidant enzymes in the embryo. It has been demonstrated that the addition of cysteamine to the *in vitro* maturation medium improves embryonic development and quality (De Matos *et al.*, 1995) and increases the intracellular GSH content of bovine oocytes (De Matos *et al.*, 1995, 2002). Generally, cysteamine doses ranging from 0.5 to 20 mM have been tested in research on semen freezing extenders and *in vitro* culture media (Balasubramanian and Rho, 2007; Bucak *et al.*, 2007; Lojkic *et al.*, 2012).

It was shown that cysteamine at a dose of 10 mM increased the number of blastocyst-stage embryos (Balasubramanian & Rhob, 2007; Lojkic *et al.*, 2012). Additionally, cysteamine at a dose of 5 mM preserved post-thaw motility at higher percentages and increased vitamin E levels after cryopreservation in ram semen (Bucak *et al.*, 2007). Hyaluronan, a nonsulphated glycosaminoglycan, is an essential component of the extracellular matrix (Erlinger, 1995). It mediates sperm functions such as sperm motility (Vines *et al.*, 2001; Ghosh *et al.*, 2002) and capacitation (Suzuki *et al.*, 2002). It has a stabilising effect on decreased motility in thawed semen (Sbracia *et al.*, 1997). Pena *et al.* (2004) successfully used hyaluronan doses of 0.5–1 mg ml<sup>-1</sup> for boar semen cryopreservation, and Uysal *et al.* (2007) used a dose of 1 mg ml<sup>-1</sup> in bull semen as a freezing extender. Fetuin is a major glycoprotein component of foetal calf serum (FCS) and a protease inhibitor, and has been shown to inhibit zona pellucida hardening during the *in vitro* maturation of mouse (Schroeder *et al.*, 1990) and equine oocytes (Dell'Aquila *et al.*, 1999). Similar to BSA, fetuin is a commercially available protein, which improves sperm motility. It was reported that at a dose of 3 mg ml<sup>-1</sup>, fetuin had no detectable effect on sperm motility, but induced forward motility at a dose of 80 mg ml<sup>-1</sup> (Jaiswal *et al.*, 2010).

The aim of this study was to compare the effectiveness of the different doses of three antioxidants including cysteamine, hyaluronan and fetuin in the freezing of Brown Swiss bull semen, and to determine the effect of these additives on sperm parameters, motility characteristics, MDA levels and SOD and CAT activities following the freeze-thawing process.

## Materials and methods

### Chemicals

The antioxidants (fetuin, cysteamine and hyaluronan) and other chemicals used in this study were obtained from Sigma–Aldrich Chemical Co. (Interlab Ltd., Ankara, Turkey).

### Animals and semen collection

Ejaculates from three Brown Swiss bulls (3 and 4 years of age) were used in the study. The bulls, belonging to the Lalahan Livestock Central Research Institute (Ankara, Turkey), were maintained under uniform feeding and housing conditions. A total number of 45 ejaculates (15 ejaculates for each bull) were collected from the bulls with the aid of an artificial vagina twice a week, according to AI standard procedures. However, 36 ejaculates (12 ejaculates for each bull) containing spermatozoa with

>80% forward progressive motility and concentrations higher than  $1.0 \times 10^9$  spermatozoa  $\text{ml}^{-1}$  were used. Immediately after collection, the ejaculates were immersed in a warm water bath at 34 °C until their assessment in the laboratory.

### Semen processing

The volume of the ejaculates was measured in a conical tube graduated at 0.1-ml intervals, and sperm concentration was determined by means of an Accucell photometer (IMV, L'Aigle, France). Sperm motility was estimated using phase-contrast microscopy (200 $\times$ ). A Tris-based extender (Trizma base 254 mM, citric acid monohydrate 78 mM, D (-) fructose 70 mM, egg yolk 15% (v/v), glycerol 6% (v/v), pH 6.8) was used as the base extender (freezing extender). Each ejaculate was split into seven equal parts and diluted to a final concentration of  $60 \times 10^6 \text{ ml}^{-1}$  spermatozoa with a Tris-based extender containing cysteamine (2.5 and 7.5 mM), hyaluronan (0.25 and 1 mg  $\text{ml}^{-1}$ ), fetuin (5 and 10 mg  $\text{ml}^{-1}$ ) and no additive (control). The cysteamine (Balasubramanian & Rhob, 2007; Bucak *et al.*, 2007; Lojkic *et al.*, 2012), hyaluronan (Pena *et al.*, 2004; Uysal *et al.*, 2007) and fetuin (Jaiswal *et al.*, 2010) doses tested in this study were determined according to previous research. Diluted semen samples were loaded into 0.25-ml French straws, cooled down to 4 °C in 2 h and frozen at a programmed rate of  $-3 \text{ }^\circ\text{C min}^{-1}$  from +4 to  $-10 \text{ }^\circ\text{C}$ ;  $-40 \text{ }^\circ\text{C min}^{-1}$  from  $-10$  to  $-100 \text{ }^\circ\text{C}$ ;  $-20 \text{ }^\circ\text{C min}^{-1}$  from  $-100$  to  $-140 \text{ }^\circ\text{C}$  in a digital freezing machine (Digitcool 5300 ZB 250; IMV). Thereafter, the straws were plunged into liquid nitrogen. After at least 24 h, the frozen straws were thawed in a 37 °C water bath for 20 s immediately before use.

### Evaluation of sperm parameters

#### *Analysis of subjective and CASA motilities*

Subjective motility was assessed using a phase-contrast microscope (100 $\times$  magnification), with a warm stage maintained at 37 °C. A wet mount was made using a 5- $\mu\text{l}$  drop of semen placed directly onto a microscope slide and covered by a cover slip. Sperm motility estimations were performed in three different microscopic fields for each semen sample. The mean of the three successive estimations was recorded as the final motility score. In addition to estimating subjective sperm motility, a computer-assisted sperm motility analysis (CASA, version 12 IVOS; Hamilton-Thorne Biosciences, Beverly, MA, USA) was also performed to analyse sperm motility characteristics. CASA was set up as follows: phase contrast; frame rate – 60 Hz; minimum contrast – 70; low and

high static size gates – 0.6 to 4.32; low and high intensity gates – 0.20 to 1.92; low and high elongation gates 7 to 91; default cell size – 10 pixels; default cell intensity – 80. Thawed semen was diluted (5  $\mu\text{l}$  semen +95  $\mu\text{l}$  extender) in a Tris-based extender (without egg yolk and glycerol) and evaluated immediately after dilution. A 4- $\mu\text{l}$  sample of diluted semen was put onto a pre-warmed chamber slide (Leja 4, Leja Products, Luzernestraat B.V., Holland, the Netherlands), and sperm motility characteristics were determined with a 10 $\times$  objective at 37 °C. The following motility values were recorded: motility (%), progressive motility (%), total sperm motility, VAP (average path velocity,  $\mu\text{m s}^{-1}$ ), VSL (straight linear velocity,  $\mu\text{m s}^{-1}$ ), VCL (curvilinear velocity,  $\mu\text{m s}^{-1}$ ), ALH (amplitude of lateral head displacement,  $\mu\text{m}$ ), STR (straightness, %) and LIN (linearity index, %). For each evaluation, 10 microscopic fields including at least 300 cells were analysed.

#### *Assessment of sperm abnormalities*

For the assessment of sperm abnormalities, of each sample, at least three drops were added into Eppendorf tubes containing 1 ml of Hancock solution [62.5 ml formalin (37%), 150 ml sodium saline solution, 150 ml buffer solution and 500 ml double-distilled water] (Schaffer & Holzmann, 2000). One drop of this mixture was put onto a slide and covered with a cover slip. The percentages of sperm acrosome and total abnormalities were determined by counting a total of 200 spermatozoa by phase-contrast microscopy (magnification 1000 $\times$ , oil immersion).

### Hypo-osmotic swelling test

The hypo-osmotic swelling test (HOST) was used to evaluate the functional integrity of the sperm membrane, based on curled and swollen tails. This was performed by incubating 30  $\mu\text{l}$  of semen with 300  $\mu\text{l}$  of a 100 mOsm hypo-osmotic solution (9 g fructose + 4.9 g sodium citrate per litre of distilled water) at 37 °C for 60 min. After incubation, 0.2 ml of the mixture was spread with a cover slip on a warm slide. Two hundred spermatozoa were evaluated (magnification 1000 $\times$ ) with bright-field microscopy. Spermatozoa with swollen or coiled tails were recorded (Revell & Mrode, 1994; Buckett *et al.*, 1997).

### Assessment of sperm DNA damage

Diluted semen samples were centrifuged at 300 g for 10 min at 4 °C. The seminal plasma was discarded, and the remaining sperm cells were washed with ( $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  free) PBS to yield a concentration of  $1 \times 10^5$  spermatozoa  $\text{cm}^{-3}$  (Arabi, 2005).

Sperm DNA damage was investigated using the single cell gel electrophoresis (comet) assay, which is generally performed under high alkaline conditions. Firstly, each microscope slide was pre-coated with a layer of 1% normal melting point agarose in PBS and dried at room temperature. Next, 100  $\mu\text{l}$  of 0.7% low melting point agarose at 37 °C was mixed with 10  $\mu\text{l}$  of the cell suspension and poured on top of the first layer. Slides were allowed to solidify for 5 min at 4 °C in a moist box. The cover slips were removed, and the slides were immersed in freshly prepared cold lysis buffer containing 2.5 M NaCl, 100 mM Na<sub>2</sub>-EDTA, 10 mM Tris, 1% Triton X-100, 10% DMSO and 40 mM dithiothreitol (pH 10) for 1 h at 4 °C. Then, the slides were incubated overnight at 37 °C in 100  $\mu\text{g ml}^{-1}$  proteinase K (Sigma-Aldrich Co., St. Louis, MO, USA) and placed in lysis buffer. The slides were removed from the lysis buffer, drained and placed in a horizontal electrophoresis unit filled with fresh alkaline electrophoresis solution, containing 300 mM NaOH and 1 mM EDTA, (pH 13), for 20 min to allow the DNA to unwind. Electrophoresis was performed for 20 min at 8 °C at 12 V and was adjusted to 250 mA. Subsequently, the slides were washed with a neutralising solution of 0.4 M Tris, pH (7.5), to remove the alkali and detergents. After neutralisation, the slides were stained with 50  $\mu\text{l}$  of 2  $\mu\text{l ml}^{-1}$  ethidium bromide and covered with a cover slip. All steps were performed under dim light to prevent further DNA damage (Haines *et al.*, 1998; Singh & Stephens, 1998). The images of 50 randomly chosen nuclei were analysed by CASP (critical assessment of protein structure prediction) (Labbe *et al.*, 2001).

Observations were made at a magnification of 400 $\times$  using a fluorescent microscope (Olympus, BX51, Tokyo, Japan). Damage was detected by a tail of fragmented DNA that migrated from the sperm head, causing a 'comet' pattern, whereas whole sperm heads, without a comet, were not considered to be damaged.

### Biochemical assays

Superoxide dismutase, GPx activities and MDA levels were measured in the samples on a UV-VIS Recording Spectrophotometer (UV-2100S; Shimadzu Co., Kyoto, Japan).

Superoxide dismutase activity was measured using the method described by Fitzgerald *et al.* (1992). Briefly, 50- $\mu\text{l}$  samples were mixed with 825  $\mu\text{l}$  of substrate solution containing 0.05 mM xanthine sodium and 0.025 mM 2-(4-iodophenyl)-3-(4-nitrophenol)-5-phenyltetrazolium chloride (INT) in a buffer solution containing 50 mM CAPS (3-(cyclohexylaminol)-1-propanesulphonic acid) and 0.094 mM EDTA (pH 10.2). One hundred and twenty-five micro litre xanthine oxidase (80 U l<sup>-1</sup>) was

added to the mixture, and the increase in absorbance was followed at 505 nm for 3 min. SOD activity was expressed as U ml<sup>-1</sup>.

Glutathione peroxidase activity was measured using the method described by Pleban *et al.* (1982). Briefly, a reaction mixture containing 1 mM Na<sub>2</sub>EDTA, 2 mM reduced glutathione, 0.2 mM NADPH, 4 mM sodium azide and 1000 U glutathione reductase in 50 mM TRIS buffer (pH 7.6) was prepared. Twenty micro litre of the samples and 980  $\mu\text{l}$  of the reaction mixture were mixed and incubated for 5 min at 37 °C. The reaction was initiated by adding 8.8 mM of hydrogen peroxide, and the decrease in absorbance was recorded at 340 nm for 3 min. GPx activity was expressed as U ml<sup>-1</sup>.

Malondialdehyde level was estimated by the measurement of thiobarbituric acid-reactive substances (TBARS) in the samples by the method described by Richard *et al.* (1991). After the reaction of MDA with thiobarbituric acid, the reaction product was followed spectrophotometrically at 532 nm, using tetramethoxypropane as a standard. The results were expressed as nmol ml<sup>-1</sup>.

### Statistical analysis

The study was replicated six times. Results were expressed as the mean  $\pm$  SEM. Sperm motility, motility characteristics, abnormalities and biochemical data were analysed by analysis of variance, followed by Tukey's *post hoc* test to determine significant differences between the groups. The groups were compared for DNA damage, using analysis of variance and Tamhane's T2 multiple-range test as a *post hoc* test. Differences with values of  $P < 0.05$  were considered to be statistically significant. Statistical analyses were performed using the SPSS 11.5 package program (SPSS Inc., Chicago, IL, USA).

### Results

#### Sperm parameters (percentages of motility, sperm motility characteristics, acrosome and total abnormalities, plasma membrane integrity and DNA damage)

The effects of cysteamine, hyaluronan and fetuin on the sperm motility and motility characteristics and morphological abnormalities, membrane integrity and damaged DNA of frozen-thawed bull semen are presented in Tables 1 and 2. The samples cryopreserved in supplemented freezing extenders had higher percentages of CASA motility and CASA total motility compared with the control group ( $P < 0.001$ ). However, the best percentages of CASA motility and CASA total motility were achieved with 10 mg ml<sup>-1</sup> of fetuin (52.6  $\pm$  2.9% and

**Table 1** Mean ( $\pm$ SE) sperm motility and motion characteristics in semen supplemented with different antioxidants of Brown-Swiss bovine semen following freeze-thawing

Groups	CASA motility (%)	CASA progressive motility (%)	CASA total motility (%)	VAP ( $\mu\text{m s}^{-1}$ )	VSL ( $\mu\text{m s}^{-1}$ )	VCL ( $\mu\text{m s}^{-1}$ )	ALH ( $\mu\text{m}$ )	STR (%)	LIN (%)
Control	30.0 $\pm$ 1.8 <sup>d</sup>	12.1 $\pm$ 1.2 <sup>c</sup>	39.1 $\pm$ 2.2 <sup>e</sup>	121.7 $\pm$ 2.4 <sup>a</sup>	85.4 $\pm$ 1.5	214.0 $\pm$ 9.0 <sup>a</sup>	7.8 $\pm$ 0.5 <sup>a</sup>	70.3 $\pm$ 1.2 <sup>de</sup>	44.0 $\pm$ 1.7 <sup>bc</sup>
Hyaluronan 0.25 mg ml <sup>-1</sup>	42.1 $\pm$ 1.8b <sup>c</sup>	13.7 $\pm$ 2.3 <sup>bc</sup>	55.4 $\pm$ 1.4 <sup>cd</sup>	119.2 $\pm$ 5.5 <sup>a</sup>	87.3 $\pm$ 4.4	204.0 $\pm$ 11.8 <sup>a</sup>	7.7 $\pm$ 0.4 <sup>a</sup>	73.9 $\pm$ 1.6 <sup>bcd</sup>	46.3 $\pm$ 1.6 <sup>bc</sup>
Hyaluronan 1 mg ml <sup>-1</sup>	40.1 $\pm$ 2.6b <sup>c</sup>	19.6 $\pm$ 1.6 <sup>a</sup>	52.7 $\pm$ 2.8 <sup>d</sup>	100.2 $\pm$ 6.5 <sup>b</sup>	78.8 $\pm$ 3.0	160.7 $\pm$ 15.4 <sup>b</sup>	6.3 $\pm$ 0.5 <sup>b</sup>	78.4 $\pm$ 1.6 <sup>a</sup>	51.1 $\pm$ 1.8 <sup>a</sup>
Fetuin 5 mg ml <sup>-1</sup>	38.1 $\pm$ 2.3 <sup>c</sup>	14.6 $\pm$ 1.4 <sup>abc</sup>	51.6 $\pm$ 1.9 <sup>d</sup>	123.5 $\pm$ 5.7 <sup>a</sup>	84.3 $\pm$ 4.1	220.7 $\pm$ 10.5 <sup>a</sup>	7.9 $\pm$ 0.4 <sup>a</sup>	68.7 $\pm$ 0.8 <sup>e</sup>	41.6 $\pm$ 1.0 <sup>c</sup>
Fetuin 10 mg ml <sup>-1</sup>	52.6 $\pm$ 2.9 <sup>a</sup>	14.0 $\pm$ 1.4 <sup>bc</sup>	64.3 $\pm$ 2.4 <sup>ab</sup>	124.3 $\pm$ 2.2 <sup>a</sup>	86.5 $\pm$ 2.7	223.5 $\pm$ 4.7 <sup>a</sup>	8.3 $\pm$ 0.3 <sup>a</sup>	71.7 $\pm$ 0.9 <sup>cde</sup>	43.3 $\pm$ 0.7 <sup>c</sup>
Cysteamine 2.5 mM	55.3 $\pm$ 2.2 <sup>a</sup>	17.4 $\pm$ 1.9 <sup>ab</sup>	69.1 $\pm$ 1.8 <sup>a</sup>	117.3 $\pm$ 3.6 <sup>a</sup>	86.6 $\pm$ 2.4	200.8 $\pm$ 9.6 <sup>a</sup>	7.8 $\pm$ 0.5 <sup>a</sup>	76.4 $\pm$ 1.9 <sup>ab</sup>	49.0 $\pm$ 2.3 <sup>ab</sup>
Cysteamine 7.5 mM	46.0 $\pm$ 2.1 <sup>b</sup>	18.1 $\pm$ 1.6 <sup>ab</sup>	59.6 $\pm$ 1.7 <sup>bc</sup>	114.1 $\pm$ 1.5 <sup>a</sup>	85.7 $\pm$ 2.3	210.9 $\pm$ 4.2 <sup>a</sup>	8.7 $\pm$ 0.3 <sup>a</sup>	75.1 $\pm$ 1.7 <sup>abc</sup>	43.9 $\pm$ 2.0 <sup>bc</sup>
P	****	*	****	**	—	****	*	****	****

\*n = 6; refers to six replications.

Different superscripts within the same column demonstrate significant differences (\* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ ).

—, no significant difference ( $P > 0.05$ ).

64.3  $\pm$  2.4%) and 2.5 mM of cysteamine (55.3  $\pm$  2.2% and 69.1  $\pm$  1.8%). The addition of 1 mg ml<sup>-1</sup> of hyaluronan (19.6  $\pm$  1.6%) and 2.5 and 7.5 mM of cysteamine (17.4  $\pm$  1.9% and 18.1  $\pm$  1.6%) provided a greater post-thaw progressive motility in comparison with the control group and the other experimental groups ( $P < 0.001$ ). For VAP, VCL and ALH, the lowest values (100.2  $\pm$  6.5%, 160.7  $\pm$  15.4% and 6.3  $\pm$  0.5%) were obtained with 1 mg ml<sup>-1</sup> of hyaluronan. For STR and LIN, the highest values (78.4  $\pm$  1.6% and 51.1  $\pm$  1.8%) were recorded for 1 mg ml<sup>-1</sup> of hyaluronan ( $P < 0.001$ ). No significant differences were observed among the groups for VSL values ( $P > 0.05$ ). In terms of sperm acrosome and total abnormalities, 10 mg ml<sup>-1</sup> of fetuin (4.3  $\pm$  0.4% and 11.7  $\pm$  0.6%) and 2.5 mM of cysteamine (4.9  $\pm$  0.3% and 13.0  $\pm$  0.7%) produced better protection levels ( $P < 0.001$ ). The results of HOST showed that the percentage values of membrane integrity in all the groups, excluding that supplemented with 5 mg ml<sup>-1</sup> of fetuin, were higher than that of the control group ( $P < 0.001$ ). As regards the DNA damage of sperm cells, 0.25 mg ml<sup>-1</sup> of hyaluronan (2.39  $\pm$  0.21%), and 2.5 and 7.5 mM of cysteamine (2.98  $\pm$  0.20% and 3.07  $\pm$  0.24%) led to lower concentrations of spermatozoa with damaged DNA than that of the control group (4.88  $\pm$  0.48%). This reduction in DNA damage was found to be statistically significant ( $P < 0.001$ ).

#### MDA levels, SOD and GPX activities

The influence of antioxidants on MDA levels and GPX and SOD activities in thawed bull spermatozoa are set out in Table 3. Supplementation with antioxidants significantly affected SOD and GPX activities in comparison with the control group ( $P < 0.001$ ). The maintenance of SOD and GPX antioxidant activities in the groups supplemented with 2.5 and 7.5 mM of cysteamine (27.60  $\pm$  5.74 U ml<sup>-1</sup> and 30.40  $\pm$  4.16 U ml<sup>-1</sup>, 3.45  $\pm$  0.65 U ml<sup>-1</sup> and 3.80  $\pm$  0.52 U ml<sup>-1</sup>, respectively) and 10 mg ml<sup>-1</sup> of fetuin (12.42  $\pm$  0.24 U ml<sup>-1</sup> and 1.56  $\pm$  0.02 U ml<sup>-1</sup>, respectively) was demonstrated to be higher than in the control group, following the freeze-thawing process ( $P < 0.001$ ). Malondialdehyde formation was found to be at a lower level in the groups supplemented with 0.25 mg ml<sup>-1</sup> of hyaluronan (0.57  $\pm$  0.02 nmol ml<sup>-1</sup>) and 7.5 mM of cysteamine (0.21  $\pm$  0.01 nmol ml<sup>-1</sup>), compared with the other experimental groups, after the freeze-thawing process ( $P < 0.001$ ).

#### Discussion

The successful cryopreservation of semen provides a high conception rate and economical benefits via artificial

**Table 2** Mean ( $\pm$ SE) sperm abnormalities, membrane integrity and DNA damage in semen supplemented with different antioxidants of Brown-Swiss bovine semen following freeze–thawing

Groups	Acrosomal abnormality (%)	Total abnormalities (%)	Membrane integrity (%)	Damaged DNA (%)
Control	11.7 $\pm$ 0.9 <sup>a</sup>	23.1 $\pm$ 0.9 <sup>a</sup>	53.0 $\pm$ 0.7 <sup>b</sup>	4.88 $\pm$ 0.48 <sup>c</sup>
Hyaluronan 0.25 mg ml <sup>-1</sup>	10.4 $\pm$ 0.5 <sup>ab</sup>	21.6 $\pm$ 1.3 <sup>ab</sup>	63.7 $\pm$ 1.1 <sup>a</sup>	2.39 $\pm$ 0.21 <sup>b</sup>
Hyaluronan 1 mg ml <sup>-1</sup>	9.6 $\pm$ 0.6 <sup>bc</sup>	22.7 $\pm$ 0.9 <sup>a</sup>	62.1 $\pm$ 1.7 <sup>a</sup>	3.47 $\pm$ 0.24 <sup>ac</sup>
Fetuin 5 mg ml <sup>-1</sup>	8.6 $\pm$ 0.6 <sup>c</sup>	19.1 $\pm$ 1.3 <sup>b</sup>	56.4 $\pm$ 1.4 <sup>b</sup>	3.59 $\pm$ 0.25 <sup>ac</sup>
Fetuin 10 mg ml <sup>-1</sup>	4.3 $\pm$ 0.4 <sup>d</sup>	11.7 $\pm$ 0.6 <sup>c</sup>	62.4 $\pm$ 2.1 <sup>a</sup>	3.43 $\pm$ 0.27 <sup>ac</sup>
Cysteamine 2.5 mM	4.9 $\pm$ 0.3 <sup>d</sup>	13.0 $\pm$ 0.7 <sup>c</sup>	63.0 $\pm$ 1.5 <sup>a</sup>	2.98 $\pm$ 0.20 <sup>ab</sup>
Cysteamine 7.5 mM	9.6 $\pm$ 0.6 <sup>bc</sup>	21.3 $\pm$ 1.1 <sup>ab</sup>	62.6 $\pm$ 1.6 <sup>a</sup>	3.07 $\pm$ 0.24 <sup>ab</sup>
<i>P</i>	***	***	***	***

'*n* = 6' refers to six replications.

Different superscripts within the same column demonstrate significant differences (\*\**P* < 0.001).

**Table 3** Mean ( $\pm$ SE) MDA levels, GPX and SOD antioxidant activities in semen supplemented with different antioxidants of Brown-Swiss bovine semen following freeze–thawing

Groups	SOD (U ml <sup>-1</sup> )	GPX (U ml <sup>-1</sup> )	MDA (nmol ml <sup>-1</sup> )
Control	2.75 $\pm$ 0.24 <sup>c</sup>	0.38 $\pm$ 0.03 <sup>c</sup>	2.31 $\pm$ 0.76 <sup>c</sup>
Hyaluronan 0.25 mg ml <sup>-1</sup>	7.88 $\pm$ 1.03 <sup>bc</sup>	0.98 $\pm$ 0.13 <sup>bc</sup>	0.57 $\pm$ 0.02 <sup>d</sup>
Hyaluronan 1 mg ml <sup>-1</sup>	4.70 $\pm$ 0.41 <sup>bc</sup>	0.61 $\pm$ 0.04 <sup>bc</sup>	1.92 $\pm$ 0.78 <sup>c</sup>
Fetuin 5 mg ml <sup>-1</sup>	8.13 $\pm$ 0.20 <sup>bc</sup>	1.02 $\pm$ 0.03 <sup>bc</sup>	2.30 $\pm$ 0.03 <sup>c</sup>
Fetuin 10 mg ml <sup>-1</sup>	12.42 $\pm$ 0.24 <sup>b</sup>	1.56 $\pm$ 0.02 <sup>b</sup>	7.26 $\pm$ 0.30 <sup>a</sup>
Cysteamine 2.5 mM	27.60 $\pm$ 5.74 <sup>a</sup>	3.45 $\pm$ 0.65 <sup>a</sup>	3.81 $\pm$ 0.12 <sup>b</sup>
Cysteamine 7.5 mM	30.40 $\pm$ 4.16 <sup>a</sup>	3.80 $\pm$ 0.52 <sup>a</sup>	0.21 $\pm$ 0.01 <sup>d</sup>
<i>P</i>	***	***	***

'*n* = 6' refers to six replications.

Different superscripts within the same column demonstrate significant differences (\*\**P* < 0.001).

insemination. Various factors including temperature changes, ice formation, oxidative stress, membrane disintegrity and damage to the acrosome and DNA reduce post-thaw semen fertility (Halliwell, 1994; Sharma & Agarwal, 1996; Thun *et al.*, 2002).

Reactive oxygen species appear to play several roles in the maintenance and disruption of spermatozoa physiology. They can also damage many cell components and induce the oxidation of cell membrane lipids, amino acids in proteins and DNA (Agarwal *et al.*, 2004). Sperm membranes are vulnerable to ROS-induced damage due to their high content of polyunsaturated fatty acids (Aurich *et al.*, 1997; Storey, 1997). Oxidative attack to sperm phospholipid-bound polyunsaturated fatty acids induces changes in membrane fluidity and integrity that impair the fertilising potential of these cells (Jones *et al.*,

1979; Aitken & Fisher, 1994). The unfavourable effects of this process can be prevented by the use of antioxidants (Sikka, 2004).

Cysteamine, a low-molecular-weight thiol compound, reacts with cystine by sulphhydryl–disulphide interchange to form a mixed disulphide (Ishii *et al.*, 1981) and enhances glutathione synthesis (Issels *et al.*, 1988; De Matos *et al.*, 2002b). Some authors reported that cysteamine supplementation of the culture medium improved blastocyst percentages (Balasubramanian & Rhob, 2007; Silva *et al.*, 2010). In the present study, the addition of 2.5 and 7.5 mM of cysteamine and 10 mg ml<sup>-1</sup> of fetuin to the semen extender provided a higher increase in SOD and GPx antioxidant enzyme activities. This was in agreement with studies performed by De Matos *et al.* (1996, 2002b), who stated that the addition of cysteamine into the maturation media resulted in an increase in GSH activity. The findings of the present study demonstrated that the addition of cysteamine to the freezing extender at concentrations of 2.5–7.5 mM could provide cryoprotection of thawed bull semen against oxidative stress-induced damage, despite the measurement of higher MDA values in the group supplemented with 2.5 mM of cysteamine. Similarly, Bucak *et al.* (2007) reported that the addition of 5 mM of cysteamine led to an increase in vitamin E levels in addition to its protective effect on sperm survival. Furthermore, Sariözkan *et al.* (2013) demonstrated that FCS had a protective role in maintaining GPx antioxidant activity and also led to a decrease in MDA levels, compared with the control group during the liquid storage of rabbit semen.

The mechanism of hyaluronan in increasing sperm motility shows that hyaluronan receptors are located in the tail, mid-piece and head of human spermatozoa (Kornovsky *et al.*, 1994). Hyaluronan supplementation may lead to an increase in phosphorylation and ATP levels, resulting in improved flagellar function and motility (Ranghanatan *et al.*, 1994, 1995; Sbracia *et al.*, 1997). The

results of the present study showed that supplementation with  $1 \text{ mg ml}^{-1}$  of hyaluronan and 2.5 and 7.5 mM of cysteamine provided a greater post-thaw progressive motility in comparison with the other groups. This observation is similar to that previously described for frozen/thawed human (Sbracia *et al.*, 1997) and mouse semen (Bakhtiari *et al.*, 2007), where hyaluronan supplementation resulted in significant increases in sperm motility and velocity movement. In contrast, Uysal *et al.* (2007) reported that hyaluronan supplementation did not provide any improvement in the post-thaw motility of bull spermatozoa. In this study, hyaluronan addition produced a significant improvement in post-thaw CASA motility. In our study, hyaluronan yielded lower VCL, VAP and ALH values in frozen/thawed bull semen. The results obtained in the present study for CASA motility are in contrast with those reported in previous research (Pena *et al.*, 2004), indicating higher VSL and VAP and lower ALH values in boar semen. The differences between the results of the present study and previous reports may be attributed to the use of different semen extenders and animal species. In the present study, all types of supplementation of the freezing extender increased CASA motility and CASA total motility, compared with the control group ( $P < 0.001$ ). Moreover,  $10 \text{ mg ml}^{-1}$  of fetuin and 2.5 mM of cysteamine had significant positive effects on the maintenance of CASA motility and CASA total motility of post-thaw bull spermatozoa when compared to the other groups.

Fetuin is an important ingredient of FCS, which is a constituent of most media used for the culture of animal cells. FCS is composed of a variety of proteins, which maintain cultured cells in a medium (Hayman *et al.*, 1985). FCS has been widely used in the culture of cumulus–oocyte complexes, because it is believed that FCS stabilises the expanding cumulus extracellular matrix (Simon *et al.*, 1992). It has been reported that fetuin inhibits zona pellucida hardening and allows normal fertilisation rates in oocytes cultured in the absence of serum (Schroeder *et al.*, 1990; Dell'Aquila *et al.*, 1999). Fetuin addition at concentrations of 5–10  $\text{mg ml}^{-1}$  and cysteamine supplementation at a dose of 2.5 mM significantly decreased acrosomal and total morphological damage of frozen-thawed bull semen, in comparison with the controls ( $P < 0.001$ ). Similarly, Sariözkan *et al.* (2013) indicated that FCS in the semen extender protected both the plasma membrane and the acrosomal integrity of chilled rabbit spermatozoa as well as sperm motility. In the present study, supplementation with  $10 \text{ mg ml}^{-1}$  of fetuin also provided for a significantly higher percentage of CASA motility, compared with the control group. Similarly, the protective effect of fetuin on motility was shown by Jaiswal *et al.* (2010). Furthermore, in the present

study, all the supplementations tested, except for fetuin at  $5 \text{ mg ml}^{-1}$ , protected the functional membrane integrity of frozen/thawed bull semen.

Apoptosis induces sperm DNA damage during cryopreservation and the thawing process (Said *et al.*, 2010). However, Thomson *et al.* (2009) suggested that increased sperm DNA damage during the freezing/thawing process results principally from oxidative stress rather than apoptosis. The Comet assay is a widely applied technique for measuring and analysing DNA breakage in individual cells (Ostling & Johanson, 1984). It has also been proven as an effective technique in evaluating the capacity of antioxidants to protect genetic material integrity in biological studies (Heaton *et al.*, 2002; Novotna *et al.*, 2007; Tuncer *et al.*, 2010). Furthermore, some authors suggest that sperm DNA integrity is a more objective marker of sperm function as opposed to sperm parameters such as motility (Twigg *et al.*, 1998b; Rajesh *et al.*, 2002).

In this study, supplementation of the semen extender with  $0.25 \text{ mg ml}^{-1}$  of hyaluronan and 2.5 and 7.5 mM of cysteamine maintained DNA integrity at a statistically significant level, compared with the control group. These results were contrary to those reported in a previous study on bovine spermatozoa (Bucak *et al.*, 2010b), indicating the protection of DNA integrity. Fetuin concentrations of 5–10  $\text{mg ml}^{-1}$  did not provide statistically higher sperm DNA integrity levels than that of the control group. In a previous report, it was suggested that supplementation with FCS provided a significant reduction in DNA damage during the liquid storage of rabbit semen, compared with the control group (Sariözkan *et al.*, 2013).

## Conclusions

The results of this study clearly suggest that all the additives used had positive effects on the maintenance of sperm motility, and the integrities of the sperm plasma membrane, morphology, and DNA, compared with the control group. Furthermore, all the supplementations tested in the present study showed strong cryoprotective effect against cryopreservation/thawing damage induced by LPO. In particular, higher SOD and GPx antioxidant activities were achieved with the addition of 2.5–7.5 mM of cysteamine. Therefore, it can be stated that the addition of antioxidants to the freezing extender could improve post-thaw sperm parameters and may be recommended to facilitate the enhancement of sperm cryopreservation techniques.

## References

- Agarwal A, Saleh RA, Bedaiwy MA (2003) Role of reactive oxygen species in the pathophysiology of human reproduction. *Fertil Steril* 79:829–843.

- Agarwal A, Nallella KP, Allamaneni SS, Said TM (2004) Role of antioxidants in treatment of male infertility: an overview of the literature. *Reprod Biomed Online* 8:616–627.
- Agarwal A, Makker K, Sharma R (2008) Clinical relevance of oxidative stress in male factor infertility: an update. *Am J Reprod Immunol* 59:2–11.
- Aitken R, Fisher H (1994) Reactive oxygen species generation by human spermatozoa. *BioEssays* 16:259–268.
- Aitken RJ, Irvine DS, Wu FC (1991) Prospective analysis of sperm–oocyte fusion and reactive oxygen species generation as criteria for the diagnosis of infertility. *Am J Obstet Gynaecol* 164:542–551.
- Aitken RJ, Gordon E, Harkiss D, Twigg JP, Milne P, Jennigs Z, Irvine DS (1998) Relative impact of oxidative stress on the functional competence and genomic integrity of human spermatozoa. *Biol Reprod* 59:1037–1046.
- Arabi M (2005) Bull spermatozoa under mercury stress. *Reprod Domest Anim* 40:454–459.
- Aurich JE, Schonherr U, Hoppe H, Aurich C (1997) Effect of antioxidants on motility and membrane integrity of chilled-stored stallion semen. *Theriogenology* 48:185–192.
- Bailey JL, Bilodeau JF, Cormier N (2000) Semen cryopreservation in domestic animals: a damaging and capacitating phenomenon. *J Androl* 21:1–7.
- Bakhtiari M, Sobhani A, Akbari M, Pasbakhsh P, Abbasi M, Hedayatpoor A, Amidi F, Sargolzaei F (2007) The effect of hyaluronic acid on motility, vitality and fertilization capability of mouse sperms after cryopreservation. *Iran J Reprod Med* 5:45–50.
- Balasubramanian S, Rho GJ (2007) Effect of cysteamine supplementation of *in vitro* matured bovine oocytes on chilling sensitivity and development of embryos. *Anim Reprod Sci* 98:282–292.
- Ball BA, Gravance CG, Medina V, Baumber J, Liu IKM (2000) Catalase activity in equine semen. *Am J Vet Res* 61:1026–1030.
- Baumber J, Ball BA (2005) Determination of glutathione peroxidase and superoxide dismutase-like activities in equine spermatozoa, seminal plasma, and reproductive tissues. *Am J Vet Res* 66:1415–1419.
- Bilodeau JF, Blanchette S, Gagnon IC, Sirard MA (2001) Thiols prevent H<sub>2</sub>O<sub>2</sub>-mediated loss of sperm motility in cryopreserved bull semen. *Theriogenology* 56:275–286.
- Bucak MN, Ateşşahin A, Varışlı O, Yuce A, Tekin N, Akçay A (2007) The influence of trehalose, taurine, cysteamine and hyaluronan on ram semen microscopic and oxidative stress parameters after freeze–thawing process. *Theriogenology* 67:1060–1067.
- Bucak MN, Sariözkan S, Tuncer PB, Sakin F, Ateşşahin A, Kulaksız R, Çevik M (2010a) The effect of antioxidants on post-thawed Angora goat (*Capra hircus ancyrensis*) sperm parameters, lipid peroxidation and antioxidant activities. *Small Ruminant Res* 89:24–30.
- Bucak MN, Tuncer PB, Sariözkan S, Başpınar N, Taşpınar M, Cayan K, Bilgili A, Akalın PP, Buyukleblebici S, Aydos S, Ilgaz S, Sunguroğlu A, Oztuna D (2010b) Effects of antioxidants on post-thawed bovine sperm and oxidative stress parameters: antioxidants protect DNA integrity against cryodamage. *Cryobiology* 61:248–253.
- Buckett WM, Luckas MJM, Aird IA, Farquharson RG, Kingsland CR, Lewis-Jones DI (1997) The hypo-osmotic swelling test in recurrent miscarriage. *Fertil Steril* 68:506–509.
- De Lamirande E, Gagnon C (1993) Human sperm hyperactivation in whole semen and its association with low superoxide scavenging capacity in seminal plasma. *Fertil Steril* 59:1291–1295.
- De Lamirande E, Leclerc P, Gagnon C (1997) Capacitation as a regulatory event that primes spermatozoa for the acrosome reaction and fertilization. *Mol Hum Reprod* 3:175–194.
- De Matos DG, Furuns CC, Moses DF, Baldassarre H (1995) Effect of cysteamine on glutathione level and developmental capacity of bovine oocyte matured *in vitro*. *Mol Reprod Dev* 42:432–436.
- De Matos DG, Furnus CC, Moses DF, Martinez AG, Matkovic M (1996) Stimulation of GSH synthesis of *in vitro* matured bovine oocytes and its effect on embryo development and freezability. *Mol Reprod Dev* 54:451–457.
- De Matos DG, Herrera C, Cortvrintd R, Smitz J, Van Soom A, Nogueira D, Pasqualini RS (2002a) Cysteamine supplementation during *in vitro* maturation and embryo culture: a useful tool for increasing the efficiency of bovine *in vitro* embryo production. *Mol Reprod Dev* 62:203–209.
- De Matos DG, Gasparini B, Pasqualini SR, Thompson JG (2002b) Effect of glutathione synthesis stimulation during *in vitro* maturation of ovine oocytes on embryo development and intracellular peroxide content. *Theriogenology* 57:1443–1451.
- Dell'Aquila ME, De Felici M, Massari S, Maritato F, Minoia P (1999) Effects of fetuin on zona pellucida hardening and fertilizability of equine oocytes matured *in vitro*. *Biol Reprod* 61:533–540.
- Desai N, Sharma R, Maker K, Sabnegh E, Agarwal A (2009) Physiological and pathological levels of reactive oxygen species in neat semen of infertile men. *Fertil Steril* 92:1626–1631.
- Dizdaroglu M, Nackerdien Z, Chao BC, Gajewski E, Rao G (1991) Chemical nature of *in vivo* DNA base damage in hydrogen peroxide-treated mammalian cells. *Arch Biochem Biophys* 285:388–390.
- Erlinger R (1995) Glycosaminoglycans in porcine lung: an ultrastructural study using cupromeromic blue. *Cell Tissue Res* 281:473–483.
- Fitzgerald SP, Campbell JJ, Lamont JV (1992) *The establishment of reference ranges for selenium. The selenoenzyme glutathione peroxidase and the metalloenzyme superoxide dismutase in blood fractions.* The fifth international symposium on selenium biology and medicine. Tennessee, USA, pp 20–23.

- Garrido N, Meseguer M, Simon C, Pellicer A, Remohi J (2004) Pro-oxidative and anti-oxidative imbalance in human semen and its relation with male fertility. *Asian J Androl* 6:59–65.
- Geva E, Lessing JB, Bartoov B, Lerner-Geva L, Zabludovsky N, Amit A (1996) The effect of antioxidant treatment on human spermatozoa fertilization rate in an *in vitro* fertilization program. *Fertil Steril* 66:430–434.
- Ghosh I, Bharadwaj A, Datta K (2002) Reduction in the level of hyaluronan binding protein (HABP1) is associated with loss of sperm motility. *J Reprod Immunol* 53:45–54.
- Gonçalves F, Barretto LSS, Arruda RP, Perri SHV, Mingoti GZ (2010) Effect of antioxidants during bovine *in vitro* fertilization procedures on spermatozoa and embryo development. *Reprod Domest Anim* 45:129–135.
- Haines G, Marples B, Daniel P, Morris I (1998) DNA damage in human and mouse spermatozoa after *in vitro*-irradiation assessed by the comet assay. *Adv Exp Med Biol* 444:789–791.
- Halliwell B (1994) Free radicals antioxidants and human disease: curiosity, cause or consequence. *Lancet* 344:721–724.
- Hayman EG, Pierschbacher MD, Suzuki S, Ruoslahti E (1985) Vitronectin—A major cell attachment-promoting protein in fetal bovine serum. *Exp Cell Res* 160:245–258.
- Heaton PR, Reed CF, Mann SJ, Ransley R, Stevenson J, Charlton CJ, Smith BH, Harper EJ, Rawlings JM (2002) Role of dietary antioxidants to protect against DNA damage in adult dogs. *J Nutr* 132:1720–1724.
- Hughes CM, Lewis SEM, McKelvey-Martin VJ, Thompson W (1996) A comparison of baseline and induced DNA damage in human spermatozoa from fertile and infertile men, using a modified comet assay. *Mol Hum Reprod* 2:613–619.
- Hughes CM, McKelvey-Martin VJ, Lewis SEM (1999) Human sperm DNA integrity assessed by the Comet and ELISA Assays. *Mutagenesis* 14:71–75.
- Ishii T, Hishimura I, Bannai S, Sugita Y (1981) Mechanism of growth promotion of L 1210 by 2-mercaptoethanol *in vitro*. *J Biol Chem* 256:12387–12392.
- Issels RD, Nagele A, Eckert KG, Wilmanns W (1988) Promotion of cystine uptake and its utilization for glutathione biosynthesis induced by cysteamine and *N*-acetyl-cysteine. *Biochem Pharmacol* 37:881–888.
- Jaiswal BS, Das K, Saha S, Dungdung SR, Majumder GC (2010) Purification and characterization of a motility initiating protein from caprine epididymal plasma. *J Cell Physiol* 222:254–263.
- Jaruga P, Zastawny TH, Skokowski J, Dizdaroğlu M, Olinski R (1994) Oxidative DNA base damage and antioxidant enzyme activities in human lung cancer. *FEBS Lett* 341:59–64.
- Jones R, Mann T, Sherins RJ (1979) Peroxidative breakdown of phospholipids in human spermatozoa: spermicidal effects of fatty acid peroxides and protective action of seminal plasma. *Fertil Steril* 31:531–537.
- Kornovsky BS, McCoshen JM, Krendenter J, Turley E (1994) The regulation of sperm motility by a novel hyaluronan receptor. *Fertil Steril* 61:935–940.
- Labbe C, Martoriati A, Devaux A, Maisse G (2001) Effect of sperm cryopreservation on sperm DNA stability and progeny development in rainbow trout. *Mol Reprod Dev* 60:397–404.
- Lojic M, Getz I, Samardzija M, Matkovic M, Bacic G, Karadjole T, Macesiz N, Folnozic I, Spoljaric B (2012) Effect of cysteamine supplementation during *in vitro* culture of early stage bovine embryos on blastocyst rate and quality. *Acta Vet Brno* 81:229–234.
- Medeiros CM, Forell F, Oliveira AT, Rodrigues JL (2002) Current status of sperm cryopreservation: why isn't it better? *Theriogenology* 57:327–344.
- Novotna B, Topinka J, Solansky I, Chvatalova I, Lnenickova Z, Sram RJ (2007) Impact of air pollution and genotype variability on DNA damage in Prague policemen. *Toxicol Lett* 172:37–47.
- Ostling O, Johanson KJ (1984) Microelectrophoretic study of radiation induced DNA damages in individual mammalian cells. *Biochem Biophys Res Commun* 123:291–298.
- Pena FJ, Johannisson A, Wallgren M, Rodriguez-Martinez H (2004) Effect of hyaluronan supplementation on boar sperm motility and membrane lipid architecture status after cryopreservation. *Theriogenology* 61:63–70.
- Pleban PA, Munyani A, Beachum J (1982) Determination of selenium concentration and glutathione peroxidase activity in plasma and erythrocytes. *Clin Chem* 28:311–316.
- Potts RJ, Notarianni LJ, Jefferies TM (2000) Seminal plasma reduces exogenous oxidative damage to human sperm, determined by the measurement of DNA strand breaks and lipid peroxidation. *Mutat Res* 447:249–256.
- Rajesh KT, Doreswamy K, Shrilatha B, Muralidhara M (2002) Oxidative stress associated DNA damage in testis of mice: induction of abnormal sperms and effects on fertility. *Mutat Res* 513:103–111.
- Ranganathan S, Ganguly AK, Datta K (1994) Evidence for presence of hyaluronan binding protein on spermatozoa and its possible involvement in sperm function. *Mol Reprod Dev* 38:69–76.
- Ranganathan S, Bharadwaj A, Datta K (1995) Hyaluronan mediates sperm motility by enhancing phosphorylation of proteins including hyaluronan binding protein. *Cell Mol Biol Res* 41:467–476.
- Revell SG, Mrode RA (1994) An osmotic resistance test for bovine semen. *Anim Reprod Sci* 36:77–86.
- Richard MJ, Arnaud J, Jurkovicz C, Hachache T, Meftahi H, Laporte F, Foret M, Favier A, Cordonnier D (1991) Trace elements and lipid peroxidation abnormalities in patients with chronic renal failure. *Nephron* 57:10–15.
- Said TM, Gaglani A, Agarwal A (2010) Implication of apoptosis in sperm cryoinjury. *Reprod Biomed Online* 21:456–462.

- Sakkas D, Urner F, Bizzaro D, Manicardi G, Bianchi PG, Shoukir Y, Campana A (1998) Sperm nuclear DNA damage and altered chromatin structure: effect on fertilization and embryo development. *Hum Reprod* 13:11–19.
- Sariozkan S, Tuncer PB, Bucak MN, Ulutaş PA (2009) Influence of various antioxidants on microscopic-oxidative stress indicators and fertilizing ability of frozen-thawed bull semen. *Acta Vet Brno* 78:463–469.
- Sariözkan S, Türk G, Cantürk F, Yay A, Eken A, Akçay A (2013) The effect of bovine serum albumin and fetal calf serum on sperm quality, DNA fragmentation and lipid peroxidation of the liquid stored rabbit semen. *Cryobiology* 67:1–6.
- Sbracia M, Grasso J, Sayme N, Stronk J, Huszar G (1997) Hyaluronic acid substantially increases the retention of motility in cryopreserved thawed human spermatozoa. *Hum Reprod* 12:1949–1954.
- Schafer S, Holzmann A (2000) The use of transmigrating and spermac stain to evaluate epididymal cat spermatozoa. *Anim Reprod Sci* 59:201–211.
- Schroeder AC, Schultz RM, Kopf GS, Taylor FR, Becker RB, Eppig JJ (1990) Fetuin inhibits zona pellucida hardening and conversion of ZP2 to ZP2f during spontaneous mouse oocyte maturation *in vitro* in the absence of serum. *Biol Reprod* 43:891–897.
- Sharma RK, Agarwal A (1996) Role of reactive oxygen species in male infertility. *Urology* 48:835–850.
- Sikka SC (2004) Role of oxidative stress and antioxidants in andrology and assisted reproductive technology. *J Androl* 25:5–18.
- Silva DS, Pereira LP, Navarro RB, Rosa DC, Pessoa GA, Silva CAM, Rubin MIB (2010) *In vitro* production of *Bos taurus indicus* embryos with cysteamine. *Anim Reprod* 7:29–34.
- Simon CL, Mao SJT, Larsen WJ (1992) Identification of a factor in fetal bovine serum that stabilizes the cumulus extracellular matrix. *J Biol Chem* 267:12380–12386.
- Singh NP, Stephens RE (1998) X-ray induced DNA double-strand breaks in human sperm. *Mutagenesis* 13:75–79.
- Storey BT (1997) Biochemistry of the induction and prevention of lipoperoxidative damage in human spermatozoa. *Mol Hum Reprod* 3:203–213.
- Suzuki K, Asano A, Eriksson B, Niwa K, Nagai T, Rodriguez Martinez H (2002) Capacitation status and *in vitro* fertility of boar spermatozoa: effects of seminal plasma. *Int J Androl* 25:84–93.
- Thomson LK, Fleming SD, Aitken RJ, De Iuliis GN, Zieschang JA, Clark AM (2009) Cryopreservation-induced human sperm DNA damage is predominantly mediated by oxidative stress rather than apoptosis. *Hum Reprod* 24:2061–2070.
- Thun R, Hurtado M, Janett F (2002) Comparison of Biociphos-Plus® and TRIS-egg yolk extender for cryopreservation of bull semen. *Theriogenology* 57:1087–1094.
- Tuncer PB, Bucak MN, Büyükleblebici S, Sariözkan S, Yeni D, Eken A, Akalın PP, Kinet H, Avdatek F, Gündoğan M (2010) The effect of cysteine and glutathione on sperm and oxidative stress parameters of post-thawed bull semen. *Cryobiology* 61:303–307.
- Twigg JP, Irvine DS, Aitken RJ (1998a) Oxidative damage to DNA in human spermatozoa does not preclude pronucleus formation at intracytoplasmic sperm injection. *Hum Reprod* 13:1864–1871.
- Twigg J, Fulton N, Gomez E, Irvine DS, Aitken RJ (1998b) Analysis of the impact of intracellular reactive oxygen species generation on the structural and functional integrity of human spermatozoa: lipid per-oxidation DNA fragmentation and effectiveness of antioxidants. *Hum Reprod* 13:1429–1436.
- Uysal O, Bucak MN, Yavaş I, Varisli O (2007) Effect of various antioxidants on the quality of frozen-thawed bull semen. *J Anim Vet Adv* 6:1362–1366.
- Vernet P, Faure J, Dufaure JP, Drevet JR (1997) Tissue and developmental distribution, dependence upon testicular factors and attachment to spermatozoa of GPX5, a murine epididymis-specific glutathione peroxidase. *Mol Reprod Dev* 47:87–98.
- Vines CA, Li MW, Deng X, Yudin AI, Cherr GN, Overstreet JW (2001) Identification of hyaluronic acid (HA) binding domain in the PH-20 protein that may function in cell signaling. *Mol Reprod Dev* 60:542–545.
- Vishwanath R, Shannon P (2000) Storage of bovine semen in liquid and frozen state. *Anim Reprod Sci* 62:23–53.
- Watson PF (1995) Recent developments and concepts in the cryopreservation of spermatozoa and the assessment of their post-thawing function. *Reprod Fertil Dev* 7:871–891.